

# Index

Page references followed by *f* denote figures; those followed by *t* denote tables.

## A

- Abdominal cavity, opening and locating female reproductive organs in, 137–138, 137f, 138f
- Accutase, 330, 533
- Acetaminophen, 199
- Acidic Tyrode's solution for removing zona pellucidae from preimplantation embryos (recipe), 780
- Acids  
disposal of, 791  
safe handling of, 790, 793
- Acid Tyrode's solution, 312–313, 536–537, 537f
- Acrosin, 34
- Acrosomal reaction, 34
- Acrosome, 34
- Activin A stock solution (10 µg/mL) (recipe), 386
- Adenoviral vectors, 462
- Adhesive, tissue, 197
- Adult mouse, 74–78  
behavioral mutants, 78  
coat color, 74–77, 74f, 75t, 76f  
germline-competent stem cells derived from, 403–430  
overview, 403–409  
protocols, 410–430  
morphological mutants, 78, 78f
- Affi-Gel Blue beads, 202
- Agarose  
as DNA prep contaminant, 248  
gel electrophoresis (*see* Electrophoresis)  
recovering DNA from gels, 252t
- Agarose plug formers, 276f
- AggreWell plates, differentiating ES cells into embryoid bodies in (protocol), 370–372
- Agouti coat color, 9, 75–77, 76f, 507
- Air bubbles, in embryo transfer pipette, 202–203, 202f
- AK-cyclo(RGDFC), 395, 396
- Albino, 3, 8, 76f, 77, 507–508
- Alcian Blue/Alizarin Red staining of cartilage and bone, 690f, 693, 760–761
- Alcian Blue staining of fetal cartilaginous skeleton, 690f, 693, 759
- Alcohol, dehydrating tissue in, 685, 744
- Alizarin Red staining of postnatal bone, 693, 762
- Alkaline phosphatase  
buffers (recipes), 717, 739, 752, 780  
staining, 50  
for activity (protocol), 689f, 691, 751–752  
for testing pluripotency and germline competence of induced pluripotent stem (iPS) cells, 408, 408f  
staining solution (recipe), 780
- Allantois, 72, 73, 73f
- Amaya Nucleofector technology, 462
- American Type Culture Collection (ATCC), 793
- AmeriClear Histology Clearing Solvent, 685
- AMH (anti-Müllerian hormone), 60–61
- Amino acids, in embryo culture media, 114, 115t
- Amnion  
development of, 68, 69f, 72  
isolation of, 151
- Analgnesia, 198–199, 199t. *See also* Analgesics; *specific protocols*
- Analgesics, 198–199  
buprenorphine, 198–199, 199t  
carprofen, 199, 199t  
flunixin meglumine, 199, 199t  
ketoprofen, 199, 199t  
meloxicam, 199, 199t
- Androgenotes, 571
- Anesthesia, 197–198. *See also* Anesthetics; *specific protocols*
- Anesthetics  
inhalation, 198  
isoflurane, 198  
ketamine/xylazine, 198, 198t  
tribromoethanol, 197–198
- Aneuploidy in ES cells, 332–333
- Animal identification, 94–100  
ear punching or notching, 98, 99f  
ear tags, 99–100, 100f  
microchip implants, 100  
suppliers of identification systems, 98t  
tattooing, 98  
toe clipping, 98
- Animals, humane treatment of, 791
- Antibiotics for selection, 469t
- Antigen II, 7
- Anti-Müllerian hormone (AMH), 60–61
- AP (alkaline phosphatase). *See* Alkaline phosphatase
- APC (*adenomatous polyposis colon*), 16
- Arms of homology, 433–434, 434f
- Artificial insemination, 597–598, 615–616, 616f, 619–622, 620f  
estrous cycle stage determination by vaginal smear (protocol), 617–618, 618f
- Artificial insemination, 597–598, 601–630  
artificial insemination, 597–598, 615–616, 616f, 619–622, 620f  
nonsurgical (protocol), 615–616, 616f  
overview, 597–598  
surgical (protocol), 619–622, 620f  
intracytoplasmic sperm injection (ICSI), 598, 623–630, 626f, 627f  
overview, 598  
protocol, 623–630, 626f, 627f  
ovary transplantation, 596, 601–603, 602f  
overview, 596  
protocol, 601–603, 602f  
subzonal insemination, 598  
in vitro fertilization, 596–597, 604–614  
discussion, 611–612  
Jackson Laboratory method, 605–608  
materials, 604–605  
Nakagata method, 608–610  
overview, 596–597  
protocol, 604–614  
recipes, 612–614  
troubleshooting, 611
- Astroglial cells, differentiation of RGLs into, 399
- ATCC (American Type Culture Collection), 793
- Autoclave safety, 790
- Autoradiography (protocol), 733–734
- Avertin. *See* Tribromoethanol
- ## B
- BABB (benzyl alcohol-benzyl benzoate), 693
- Bacteria  
biological safety procedures, 792–793  
shipping requirements, 793
- Bacterial artificial chromosomes (BACs), 238, 242–243, 246–247  
DNA isolation using NucleoBond system (protocol), 271–272  
DNA purification (protocol), 273  
injection buffer preparation for BAC/YAC DNA (protocol), 290
- Bacterial cell contents, as DNA prep contaminant, 247
- Bacteriophage P1, 243
- Bailey, Donald W., 8
- Balancer chromosomes, 16
- BALB/c strain, 8, 492, 493
- Bases  
disposal of, 791  
safe handling of, 790, 793

- Basic fibroblast growth factor (bFGF), 389  
Bateson, William, 3  
BDH patillated Fibrowax, 685  
Behavioral mutants, 78  
Bennett, Dorothea, 10  
Benzyl alcohol-benzyl benzoate (BABB), 693  
 $\beta$ -actin promoter, 245, 455  
 $\beta$ -galactosidase activity, 244  
  staining for activity, 689f, 691, 745–747  
 $\beta$ -mercaptoethanol (recipe), 360  
Bevelers, 775  
bFGF (basic fibroblast growth factor), 389  
Biarmed (Robertsonian fusion) chromosomes, 9  
Biggers, John, 12  
BIO 101 GENECLEAN Spin Turbo kit, 250t  
Biological safety cabinets, 770–771  
Biological safety procedures, 792–793  
Blasticidin S, 465t  
Blastocyst  
  attachment, 39  
  chimera formation  
    disaggregating cleavage-stage embryos  
      and inner cell mass of blastocysts  
      into individual cells (protocol),  
      547–548  
    injecting blastocysts (protocol), 522–525  
    isolating inner cell mass of blastocysts  
      (protocol), 549–550, 549f  
    preparation for injection or aggregation,  
    503–505, 504f  
    transfer of chimeric blastocysts, 506–507  
  collection (protocol), 146–148, 147f  
  epiblast, 38f, 41–45, 43t, 44t  
  formation, 35–37, 37f  
  handling blastocysts for fixation (protocol),  
  700  
  implantation, 37, 39, 40f, 149  
BlastoKit, 504  
Blocking solution (recipe), 718  
Blood collection by tail bleeding (protocol),  
  230–231  
*bls* gene, 465t  
BMOC (Brinster's medium for ovum culture),  
  110, 111t  
Bone, staining of, 690f, 693–694, 760–762  
Bone morphogenetic protein (BMP), 51  
Bouin's fixative, preparation of (protocol), 699  
Boveri, Theodor, 12  
Bovine serum albumin (BSA)  
  BSA fraction V in PBS (0.1%, w/v) (recipe),  
  386  
  BSA fraction V in PBS (0.1%, w/v) (recipe),  
  377  
*Brachyury* gene, 10–11  
Bradley's Allan, 466  
Braking pipette, 222, 223, 223f  
Branchial arches, formation of, 66–67, 67f  
Breeding chimeric mice, 507–509  
Breeding colony. *See* Colony  
Brinster, Ralph, 12, 13, 14  
Brinster's medium for ovum culture (BMOC),  
  110, 111t  
Brown coat color, 76f, 77  
Buffers and solutions, 779–784. *See also* Recipes  
Buprenorphine, 198–199, 199t  
  
**C**  
CaCl<sub>2</sub> (0.1 M) (recipe), 392  
Caesarean section  
  overview, 203  
  protocol, 220–221  
Capacitation, 34  
Capecchi, Mario, 18  
Capillaries, pulling embryo-handling pipettes  
  from, 134–135, 774–775  
Carbon dioxide, for embryo culture, 116  
Carprofen, 199, 199t  
Cartilage, staining of, 690f, 693, 759–761  
Castle, William E., 3  
Castration  
  overview, 204  
  protocol, 234–235  
CAT (chloramphenicol acetyltransferase) gene,  
  244  
C-banding, 332  
C57BL/6 strains, 324, 492–493, 497f–498f,  
  497t–498t, 507–508  
  C57BL/6 J strain, 1, 508  
  C57BL/6N strain, 508  
cDNA  
  addition of heterologous introns to, 245  
  expression of, 245  
Cell fusion induced by Sendai virus, 572,  
  580–581, 581f  
Cervical dislocation, 137, 137f  
Cesium chloride, as DNA prep contaminant,  
  248  
Cesium chloride gradient centrifugation, for  
  contaminant removal in DNA  
  prep, 248  
CGH (comparative genomic hybridization), for  
  copy number variation  
  determination, 333  
Chemically defined medium (CDM) + FGF2  
  and ActA medium for culture in  
  feeder-free conditions (recipe),  
  387  
Chemically defined medium (CDM) for culture  
  in feeder-free conditions (recipe),  
  387  
Chemicals, properties of common hazardous,  
  793–794  
Chicago Sky Blue, 157  
Chicago Sky Blue 6B (Pontamine Sky Blue)  
  (recipe), 781  
Chicken  $\beta$ -actin promoter, 245, 455  
Chimeras, 13–14, 489–550  
  breeding chimeric mice, 507–509  
  color coat markers, 491–493, 494f–499f,  
  494t–500t  
  in generation of C57BL6 ES cell  
  chimeras, 497f–498f, 497t–498t  
  in generation of 129 ES cell chimeras,  
  494f–496f, 494t–496t  
  in generation of ES cell chimeras using F<sub>1</sub>  
  hybrid and albino cell lines, 499f,  
  499t–500t  
ES cell injection chimeras, 505–506  
  microinjection setup, 505–506  
  preparation of injection and holding  
  pipettes, 506  
  genotyping, 501, 501f, 502  
  overview, 490–500  
  preparation of embryos for injection or  
  aggregation, 503–505, 504f  
  protocols, 514–550  
    aggregate assembly, 542–546  
      diploid embryo ↔ diploid embryo,  
      542–543, 543f  
      diploid embryo ↔ tetraploid embryo,  
      544–545  
      ES cells ↔ diploid embryo, 543–544,  
      544f  
    ES cells ↔ tetraploid embryo,  
      545–546, 546f  
      materials, 542  
      method, 522–524, 523f, 524f  
      troubleshooting, 524–525  
    injecting blastocysts, 522–525  
      discussion, 525  
      materials, 522  
      method, 522–524, 523f, 524f  
      troubleshooting, 524–525  
    injecting eight-cell-stage embryos,  
      526–530  
      discussion, 528  
      materials, 526  
      method, 527–528  
      recipes, 529–530  
      troubleshooting, 528–529  
    isolating inner cell mass of blastocysts,  
      549–550, 549f  
    making ES cell injection needles,  
      514–517, 514f, 516f  
    tetraploid embryo production, 539–541,  
      540f, 540t  
    zona pellucida removal, 536–538, 537f  
  sex chromosomes, 493, 500  
  transfer of chimeric blastocysts, 506–507  
  troubleshooting production and breeding,  
  509–511  
  types of, 500–503, 501f, 501t  
    diploid embryo ↔ diploid embryo,  
    500–501, 501t, 542–543, 543f  
    diploid embryo ↔ tetraploid embryo,  
    501t, 502, 544–545  
    ES cells ↔ diploid embryo, 501–502,  
    501t, 543–544, 544f  
    ES cells ↔ tetraploid embryo, 501t,  
    502–503, 545–546, 546f  
Chloramphenicol acetyltransferase (CAT) gene,  
  244  
Chloroform, as DNA prep contaminant, 248  
Chorioallantoic placenta, 68, 69f, 72–74, 73f  
Chromosomal banding, 332  
Chromosome(s)  
  aberrations, induced site-specific, 449–451  
  deletions, 450  
  inversions, 450  
  translocations, 450–451  
  anomalies in ES cells, 332–333  
  balancer, 16  
  biarmed (Robertsonian fusion), 9  
  karyotyping, 332–333, 363–365, 364f  
  sex chromosomes in chimeras, 493, 500  
Chromosome counting (protocol), 363–365,  
  364f  
Circulation system, development stages, 26t–27t  
CL (corpora lutea), 32, 201  
Cleavage, 35, 36f  
*Clock*, 16  
Cloning mice  
  overview, 572–573  
  protocol, 583–590  
  collecting and enucleating oocytes,  
  587–588

- discussion, 590  
 donor nucleus cell preparation, 588  
 embryo activation and culture, 589–590  
 enucleation and injection micropipette setup, 586–587, 587f  
 injecting donor nuclei, 588–589, 589f  
 materials, 583–584, 584t, 585f, 586f  
 preparing micropipettes, 585  
 recipe, 590  
 strain characteristics of recipient oocytes, 585  
 transferring cloned embryos, 590
- Closure, of surgical incisions, 197
- CMV (cytomegalovirus) enhancer, 245, 455
- c-MYC transcription factor and gene, 406, 407
- Coat color, 74–77, 74f, 75t, 76f  
 agouti, 75–77, 76f, 507  
 albino, 3, 8, 76f, 77, 507–508  
 brown, 76f, 77  
 dilute, 9, 10, 76f, 77  
 markers in ES cell chimera formation, 491–493, 494f–499f, 494t–500t  
 in generation of C57BL6 ES cell chimeras, 497f–498f, 497t–498t  
 in generation of 129 ES cell chimeras, 494f–496f, 494t–496t  
 in generation of ES cell chimeras using F<sub>1</sub> hybrid and albino cell lines, 499f, 499t–500t  
 pink-eyed dilution, 3, 76f, 77, 507
- Colcemid, in pronuclear transfer protocol, 572, 577–582
- Collaborative Cross, 8
- Collection of preimplantation embryos, 123  
 blastocyst collection (protocol), 146–148, 147f  
 opening abdominal cavity and locating female reproductive organs (protocol), 137–138, 137f, 138f  
 pipettes for handling embryos, 121–122, 122f, 134–137  
 two-cell- to morula-stage embryo collection (protocol), 143–145, 144f–145f  
 zygote collection and removal of cumulus cells (protocol), 139–142, 140f–141f
- Colony, 85–108  
 animal identification, 94–100  
 ear punching or notching, 98, 99f  
 ear tags, 99–100, 100f  
 microchip implants, 100  
 suppliers of identification systems, 98t  
 tattooing, 98  
 toe clipping, 98  
 diet used, 94  
 embryo donors, 89–92  
 female for generation of ES cell chimeras, 90, 92  
 female for pronuclear microinjection, 89–90, 91–92, 92t  
 production, 91–92  
 stud male mice for, 91  
 embryo recipients, 92–93  
 production of, 93  
 sterile male mice to induce pseudopregnancy, 93  
 establishing breeding colony to supply embryo donor and recipient mice, 88–89  
 location of embryo manipulation laboratory, 87–88  
 natural matings, setting up, 94–95, 95t  
 pathogen control in, 86–87  
 pregnancy with manipulated embryos, 94  
 protocols, 103–108  
 intraperitoneal injection, 107–108, 107f  
 preparation of gonadotropins and superovulation, 105–106  
 selecting females in estrus and checking plugs, 103, 104f  
 superovulation induction, 95–97, 96t  
 age and weight, 96–97, 96t  
 gonadotropin dose, 96t, 97  
 time of administration, 97  
 performance of stud males, 97  
 strain background, 96, 96t
- Color. *See* Coat color
- Comparative genomic hybridization (CGH), for copy number variation determination, 333
- Complement, guinea pig serum (recipe), 360
- Complete ES cell medium (ES-DMEM) (recipe), 327t, 430
- Complete ES cell medium for C57BL6/N ES cells (recipe), 529
- Complete ES cell medium for 129S6/SvEvTac × C57BL/6N F<sub>1</sub> hybrid ES cells (recipe), 530
- Conditional genetic manipulations, 15
- Contaminants in DNA isolation and purification removal procedures, 248–249  
 cesium chloride gradient centrifugation, 248  
 dialysis, 248  
 electroelution, 248  
 ethanol precipitation, 248  
 extraction with *n*-butanol, 248  
 gelase, 248  
 heat inactivation of enzymes, 248  
 ion-exchange chromatography, 248  
 phenol:chloroform extraction, 248  
 silica matrix columns, 249  
 types/sources of, 247–248
- Copulation plug. *See* Matings; Vaginal plug
- Copy number determination, 553
- Copy number variation in ES cell lines, 333
- Corpora hemorrhagica, 201
- Corpora lutea (CL), 32, 201
- Cre/loxP* system, 239
- Cre* recombinase  
 chromosomal aberrations, induced site-specific, 449–451  
 described, 17, 239, 437–438  
 in ES cell-mediated transgenesis, 441–442, 455–456  
 in gene targeting examples, 443, 444–445  
 site-specific recombination-mediated insertions, 448–449  
 in tamoxifen- and RU486-inducible systems, 440
- CRISPR/Cas system, 1, 434, 434f
- Cryopreservation, 631–681, 777  
 embryo  
 equilibrium methods, 633  
 factors to consider, 632–633  
 by high-osmolality vitrification (protocol), 658–664, 660f, 661f  
 cooling embryos, 659–660, 662  
 discussion, 661  
 in insemination straws, 662–664  
 loading embryos into cryotube, 660f  
 materials, 658–659  
 warming embryos, 660–661, 661f, 662–663  
 nonequilibrium methods, 634–635  
 number of embryos to preserve, 633  
 by rapid cooling (protocol), 652–657  
 discussion, 656–657  
 materials, 652–653  
 method, 653–655, 655f  
 recipe, 657  
 warming embryos, 655–656  
 by slow freezing (protocol), 646–651  
 materials, 646–647  
 method, 647–649  
 recipes, 650–651  
 straw preparation and loading, 648, 648f  
 thawing embryos, 649  
 troubleshooting, 650  
 thawing embryos, 649  
 warming embryos, 655–656, 660–661, 661f, 662–663
- of epiblast stem cells (EpiSCs), 386
- of ES cells (protocol), 345–347  
 discussion, 347  
 freezing ES cells, 346  
 materials, 345  
 thawing ES cells, 346–347
- of extra-embryonic endoderm stem (XEN) cells, 381
- oocyte, 636
- ovary  
 overview, 637  
 protocol, 676–678  
 records, 637
- sperm  
 overview, 635–636  
 using cryoprotectant containing L-glutamine (protocol), 671–675, 672f–674f  
 cryopreservation, 673  
 materials, 671–672, 672f  
 recipe, 674–675  
 thawing sperm, 673–674  
 using cryoprotectant containing monothio glycerol (MTG) (protocol), 665–670  
 cryopreservation, 667  
 materials, 665–666  
 recipe, 668–669  
 setup, 666–667  
 sperm collection, 667  
 straw preparation and loading, 666, 666f  
 thawing sperm, 668  
 troubleshooting, 668–669
- of spermatogonial stem cells (protocol), 416–417
- storage of samples, 637
- transport of embryos and germ plasm, 640–641
- of trophoblast stem (TS) cells, 376
- Cryoprotectant solution, 1.5 M 1,2-propanediol (recipe), 650
- Cryoprotective agent with 100 mM L-glutamine (gCPA) (recipe), 674–675
- Cryoprotective medium (CPM) (recipe), 669
- Cuénot, Lucien, 3
- Culture media  
 for preimplantation embryos, 109–117  
 choice and experimental outcome, 118  
 commercially available media, 120t  
 composition of selected media, 111t

- Culture media (*Continued*)  
 considerations for successful culture, 118–120  
 embryo collection, 118–119  
 medium choice, 118  
 quality control, 119–120  
 reagents, 118–119  
 critical components of, 113–117, 115t  
 amino acids, 114, 115t  
 carbon dioxide, 116  
 EDTA, 116–117  
 glucose, 113–114  
 growth factors, 117  
 light exposure, 117  
 osmolarity, 115  
 oxygen tension, 115–116  
 pH, 116  
 temperature, 117  
 volume of culture, 117  
 for development from zygotes to blastocysts, 112–113  
 history, 109–110  
 for overcoming two-cell block, 109  
 preparation of, 120–121, 126–131  
 concentrated stocks, 129  
 HEPES-buffered media for embryo handling, 128–129, 128t  
 media from concentrated stocks, 129–131, 130t  
 overview, 120–121  
 protocol, 126–131, 127t, 128t, 130t  
 Culture medium (recipe), 174, 186  
 Culture of embryos. *See* Culture media;  
 Postimplantation embryos,  
 culture of; Preimplantation  
 embryos, culture of  
 Cumulina the mouse, 13, 403  
 Cumulus cells  
 removal with hyaluronidase, 141  
 sperm penetration of, 34  
 Cutting devices, safe handling of, 791  
 Cutting sections  
 cutting thick sections using a vibratome  
 (protocol), 709–711  
 fixed specimens, 710–711  
 live specimens, 712–713  
 materials, 709, 712  
 cutting wax sections using a microtome  
 (protocol), 706–707  
 overview, 685  
 transferring sections onto water droplets  
 (protocol), 708  
 Cytochalasin B, in pronuclear transfer protocol,  
 572, 577–582  
 Cytomegalovirus (CMV) enhancer, 245, 455  
 CZB media, 111t, 112, 126, 127t, 130t, 576, 584t,  
 629t–630t (recipe)  
 CZB HEPES, 128t, 130t, 584t, 629t–630t  
 (recipe)  
 CZB-Sr medium, 584t
- D**  
 DAPI staining of whole-mount embryos or fetal  
 organs (protocol), 722–723, 723f  
 DBA strain, 7  
 Decidual reaction, 39  
 Deciduum, 39  
 Dehydrating tissue in alcohol, 685, 744  
 Deletions, in ES cell lines, 332–333  
 Department of Health, Education, and Welfare  
 (HEW), U.S., 793  
 DEPC (diethyl pyrocarbonate), 685  
 DEPC-treated water (recipe), 697, 705  
 Dermomyotome, 57–58, 57f  
 Detergent rinse (recipe), 749  
 Development, 23–79  
 branchial arch and pharyngeal region  
 formation, 66–67, 67f  
 cell-autonomous markers used to follow  
 lineages, 46t  
 chorioallantoic placenta, 68, 69f, 72–74, 73  
 cleavage, 35, 36f  
 embryonic turning, 54, 55f  
 endoderm  
 anterior visceral, 52  
 formation of definitive, 45–49  
 gut, 68  
 parietal, 68–71, 69f–71f  
 primitive, 38f, 41, 42f  
 visceral, 68–71, 69f–71f  
 epiblast, 38f, 41–45, 42f, 43t, 44t  
 extra-embryonic tissues, 68–70, 69f, 70f  
 fertilization, 33f, 34–35  
 gastrulation, 45–49, 48f  
 germline origin, 49–51, 50f  
 gestation and embryo staging, 24–27, 25f,  
 26t–27t, 28f–30f  
 gut, 67–68  
 heritage of experimental embryology, 11–14  
*Hox* genes, 58, 59f  
 implantation, 37, 39, 40f  
 kidney, 60  
 lateral plate, 58, 60  
 limb formation, 61  
 lineages of tissues, 43f  
 mesoderm, 43, 43t  
 differentiation of extra-embryonic, 72,  
 72f  
 formation of, 45–49  
 generation of regional diversity in, 51–52  
 intermediate, 58, 60  
 morula compaction and blastocyst  
 formation, 35–37, 37f  
 neural crest, 65–66, 66f  
 neurulation, 61–62  
 node, 52–54, 53f  
 oocyte maturation, 27, 30, 31f–32f  
 origins of developmental genetics, 9–11  
 ovulation, 30, 32, 33f, 34f  
 pregastrula embryo, 45, 47f  
 preimplantation, 28f, 36f  
 schematic from implantation to neurula  
 stage, 38f  
 sex determination and differentiation, 60–61  
 somites and derivatives, 55–58, 56f, 57f  
 tail bud, 54  
 time course, 25f  
 trophoctoderm, 36–37, 37f, 38f, 39, 41, 42f  
 zygote to eight-cell uncompact morula, 35  
 Devitrification, 634  
 Dewaxing and rehydrating sections  
 overview, 686  
 protocol, 714  
 DIA/LIF (differentiation inhibiting activity/  
 leukemia inhibitory factor),  
 323–324  
 Dialysis, for contaminant removal in DNA prep,  
 248  
 Diapause, induction of, 39, 204, 324, 350  
 Diestrus, 95t  
 Diethyl pyrocarbonate (DEPC), 685  
 Differentiating ES cells into embryoid bodies  
 in AggreWell plates (protocol), 370–372  
 by suspension culture (protocol), 368–369  
 Differentiation inhibiting activity/leukemia  
 inhibitory factor (DIA/LIF),  
 323–324  
 Digestion buffer (recipe), 559  
 Digoxigenin, 687  
 DiI labeling, 53  
 Dilute coat color, 9, 10, 76f, 77  
 Dilute-short ear, 10  
 Dilution solution, 1 M sucrose (recipe), 651  
 Dimethyl sulfoxide (DMSO), use in somatic cell  
 nuclear transfer (SCNT), 573  
 Diphtheria toxin (dt), 436  
 Disaggregating cleavage-stage embryos and inner  
 cell mass of blastocysts into  
 individual cells (protocol), 547–548  
 Disposal  
 general cautions, 789–791  
 of laboratory waste, 791  
 Dissecting microscopes, fluorescent protein  
 visualization with, 692–693  
 Dissection medium (recipe), 186  
 Dissociation medium (recipe), 393  
 DMEM (Dulbecco's modified Eagle's medium),  
 high glucose, 327  
 DMEM/F12 + N2 medium (recipe), 360  
 DMEM/Ham's F12, 327  
 DMSO (dimethylsulfoxide) (1.5 M)  
 cryoprotectant solution (recipe),  
 677  
 DNA  
 concentration determination, 253–254  
 introduction of foreign DNA into ES cells,  
 461–487  
 overview, 462–466  
 culture conditions, 462  
 double-stranded breaks, 434–435,  
 434f  
 electroporating DNA, 464–465, 464t  
 general considerations, 462–464  
 genotyping assays, 462, 462t  
 isolating ES cell colonies, 465–466  
 purifying DNA, 464  
 selection methods, 465, 465t  
 targeting approached, 463–464, 463f  
 transfection methods, 462–463  
 protocols, 468–487  
 electroporating DNA into cells,  
 468–471, 469t  
 genotyping ES cell colonies before  
 picking, 485–487  
 isolating ES cell colonies, 472–474  
 passaging and freezing ES cells in  
 96-well plates, 475–477  
 preparation of DNA from cells in  
 24-well tissue culture plates,  
 483–484  
 rapid preparation of DNA from cells  
 in 96-well tissue culture plates,  
 480–482  
 thawing ES cells from a 96-well plate,  
 478–479  
 isolation (*see* DNA isolation and  
 purification)  
 purification (*see* DNA isolation and  
 purification)  
 storage of, 254  
 DNA isolation and purification  
 commercial kits for, 249, 250t  
 contaminants  
 removal procedures, 248–249  
 types/sources of, 247–248

- for introduction into ES cells, 464, 480–484
  - preparation of DNA from cells in 24-well tissue culture plates (protocol), 483–484
  - rapid preparation of DNA from cells in 96-well tissue culture plates (protocol), 480–482
- overview, 247–252
  - contaminant removal, 247–249
  - digestion of plasmid DNA for release of genetic construct fragment, 251
  - fragment purification, 251–252
  - gel separation, 251
  - general considerations, 247
  - preparation of standard-size plasmid-DNA from bacterial cultures, 249–251
  - recovery of DNA from gels, 251, 251t
- preparation of DNA for microinjection, 252–254
  - DNA concentration determination, 253–254
  - filtration of DNA solution, 254
  - quality issues, 252–253
  - storage of DNA preparation, 254
- protocols, 269–291
  - BAC DNA isolation using NucleoBond system, 271–272
    - materials, 271
    - method, 271–272
    - troubleshooting, 272
  - BAC DNA purification, 273
  - DNA fragment isolation and purification, 269–270
    - materials, 269
    - method, 269–270
    - troubleshooting, 270
  - DNA preparation from cultured cells for Southern blot or PCR, 567–568
  - injection buffer preparation for BAC/YAC DNA, 290
  - injection buffer preparation for standard-sized DNA, 289
  - isolation of high-molecular weight DNA from embryonic or extra-embryonic tissues, 560–561
  - isolation of high-molecular weight DNA from mouse tail tips, 558–559
  - YAC DNA purification using PFGE and ultrafiltration, 286–288
    - materials, 286–287, 287f
    - method, 287–288
    - troubleshooting, 288
  - YAC DNA purification using two-gel electrophoresis procedure, 279–285
    - discussion, 283–284
    - excision of YAC DNA, 281
    - isolation of YAC DNA, 281–283
      - materials, 279–280
      - method, 280–283
    - PFGE gel electrophoresis, 280–281, 280t, 282f
    - recipes, 284–285
    - standard agarose gel electrophoresis, 281, 282f
  - yeast agarose plug preparation for YAC DNA isolation, 274–278
    - discussion, 276
    - materials, 274, 276f
    - method, 275–276
    - recipes, 276–277
- DNA polymerase I Klenow fragment, 739
- DNase, 505
- Dobrovolskaia-Zavadskaja, Nelly, 10
- Dolly the sheep, 403
- Dominant white spotting* pigmentation mutant, 9
- Dot-blot hybridization, for copy number determination, 553
- Double-stranded DNA breaks (DSBs), 433–434, 434f
- Doxycycline-inducible reprogramming of fibroblasts, 428–430
- D-PBS medium, 111t
- Dre recombinase, 438
- DSBs (double-stranded DNA breaks), 433–434, 434f
- dsRed fluorescent protein, 437, 692
- dt (diphtheria toxin), 436
- Dulbecco's modified Eagle's medium (DMEM) with high glucose, 327
- Dulbecco's modified Eagle's medium (DMEM)/F12 + N2 medium (recipe), 360
- Dulbecco's modified Eagle's medium (DMEM)/Ham's F12, 327
- Dunn, L.C., 10
- Dye injection, visualizing early implantation sites by, 150, 150f, 157–158
- Dyes, hazards of, 793
- E**
- Eagle's medium, amino acids in, 114, 115t
- Early gastrula organizer (EGO), 54
- Ear punching or notching, 98, 99f
- Ear tags, 99–100, 100f
- EC (embryonal carcinoma) cells, 13–14, 44, 322–323
- ECFP (enhanced cyan fluorescent protein), 690f, 692
- Ectoderm, 43, 43t
- EDTA, in embryo culture media, 116–117
- Efferent duct injection, 421f, 422
- EFS42.5c vitrification solution (recipe), 663–664
- EFS20 solution, 662
- EFS40 solution, 653–654, 656–657
- EFS20 vitrification solution (recipe), 663
- EGF (epidermal growth factor), 389
- EGFP (enhanced green fluorescent protein), 244, 690f, 692
- Eight-cell-stage embryos, injecting (protocol), 526–530
- Electrodes, for electroporation, 187, 188f
- Electroelution, for contaminant removal in DNA prep, 248
- Electrofusion, in tetraploid embryo production (protocol), 539–541, 540f, 540t
- Electrofusion apparatus, 776
- Electrolysis, sharpening tungsten needles by, 160f
- Electrophoresis
  - gel separation of DNA fragments, 251
  - recovery of DNA from agarose gels, 252t
  - YAC DNA purification using PFGE and ultrafiltration (protocol), 286–288
  - YAC DNA purification using two-gel electrophoresis procedure (protocol), 279–285
- Electroporation
  - of DNA into cells
    - overview, 464–465, 464t
    - protocol, 468–471, 469t
  - integrating *piggyBac* transposon transgenes into mouse fibroblasts by electroporation (protocol), 424–425
  - of postimplantation embryos
    - overview, 154–155, 155f
    - in utero (protocol), 190–193, 192f
    - in vitro (protocol), 187–189, 188f
- Elutip-D kit, 250t
- Embedding in wax
  - fixation and paraffin embedding for GFP visualization (protocol), 757–758
  - overview, 685
  - protocol, 701–702
  - small embryos (protocol), 703
- Embryo(s), transport of, 640–641
- Embryo collection
  - culture media for, 118–119
  - preimplantation embryos, 123
    - blastocyst collection (protocol), 146–148, 147f
    - opening abdominal cavity and locating female reproductive organs (protocol), 137–138, 137f, 138f
    - pipettes for handling embryos, 121–122, 122f, 134–137
    - two-cell- to morula-stage embryo collection (protocol), 143–145, 144f–145f
    - zygote collection and removal of cumulus cells (protocol), 139–142, 140f–141f
- Embryo cryopreservation
  - equilibrium methods, 633
  - factors to consider, 632–633
  - by high-osmolality vitrification (protocol), 658–664, 660f, 661f
  - nonequilibrium methods, 634–635
  - number of embryos to preserve, 633
  - by rapid cooling (protocol), 652–657
  - by slow freezing (protocol), 646–651
  - thawing embryos, 649
  - warming embryos, 655–656, 660–661, 661f, 662–663
- Embryo dilution solution (recipe), 657
- Embryo donors, 89–92
  - establishing breeding colony to supply, 88–89
  - female for generation of ES cell chimeras, 90, 92
    - production of donors, 92
    - strains for, 90
  - female for pronuclear microinjection, 89–90, 91–92, 92t
    - production of donors, 91–92, 92t
    - strains for, 89–90
  - production, 91–92
  - stud male mice for, 91
- Embryo-handling pipette, 121–122, 122f
  - components of, 121–122, 122f
  - hand-controlled, 121–122, 122f
  - making from hard glass capillaries, 134–135
  - reducing capillary action in, 122
  - siliconized, 136
- Embryoid body formation, 333, 366–370
- Embryonal carcinoma (EC) cells, 13–14, 44, 322–323
- Embryonic ectoderm development (eed)* locus, 10
- Embryonic stem (ES) cell(s). *See also* Embryonic stem (ES) cell lines

- Embryonic stem (ES) cell(s). *See also* Embryonic stem (ES) cell lines (*Continued*)
- chimera formation
- aggregation with embryos
    - aggregation plate preparation (protocol), 534–535, 535f
    - ES cell preparation for aggregation (protocol), 531–533, 533f
    - number of cells for, 493
    - overview, 490–493
    - tetraploid embryo production (protocol), 539–541, 540f, 540t
    - zona pellucida removal (protocol), 536–538, 537f
  - cell lines for, 491–493
  - color coat markers, 491–493, 494f–499f, 494t–500t
    - in generation of C57BL6 ES cell chimeras, 497f–498f, 497t–498t
    - in generation of 129 ES cell chimeras, 494f–496f, 494t–496t
    - in generation of ES cell chimeras using F<sub>1</sub> hybrid and albino cell lines, 499f, 499t–500t
  - embryo donors for, 90, 92
    - production of donors, 92
    - strains for, 90
  - ES cells ↔ diploid embryo aggregate assembly (protocol), 543–544, 544f
  - ES cells ↔ tetraploid embryo aggregate assembly (protocol), 545–546, 546f
  - injection into blastocyst
    - assembling microinjection setup, 518–519
    - disaggregating cleavage-stage embryos and inner cell mass of blastocysts into individual cells (protocol), 547–548
    - ES cell preparation for injection, 520–521
    - injecting blastocysts (protocol), 522–525
    - injecting eight-cell-stage embryos (protocol), 526–530
    - isolating inner cell mass of blastocysts (protocol), 549–550, 549f
    - making ES cell injection needles (protocol), 514–517, 514f, 516f
    - microinjection setup, 505–506
    - number of cells to inject, 493
    - overview, 490–493
    - preparation of injection and holding pipettes, 506
  - cryopreservation of, 345–347
  - detection and analysis of ES cell-mediated gene/genome alterations, 554–555
  - developmental origin of, 44
  - DNA introduction into, 461–487
    - overview, 462–466
      - culture conditions, 462
      - electroporating DNA, 464–465, 464t
      - general considerations, 462–464
      - genotyping assays, 462, 462t
      - isolating ES cell colonies, 465–466
      - purifying DNA, 464
      - selection methods, 465, 465t
      - targeting approached, 463–464, 463f
      - transfection methods, 462–463
    - protocols, 468–487
      - electroporating DNA into cells, 468–471, 469t
      - genotyping ES cell colonies before picking, 485–487
      - isolating ES cell colonies, 472–474
      - passaging and freezing ES cells in 96-well plates, 475–477
      - preparation of DNA from cells in 24-well tissue culture plates, 483–484
      - rapid preparation of DNA from cells in 96-well tissue culture plates, 480–482
      - thawing ES cells from a 96-well plate, 478–479
      - making homozygous for mutations, 451–452
- Embryonic stem (ES) cell lines, 321–404
- derivation and culture, overview of, 322–330
    - considerations for optimal ES cell culture, 325–327, 326f
    - historical background, 322–325
    - leukemia inhibitory factor (LIF), 329–330
    - media and reagents, 327–328, 327t, 328t
    - serum and serum replacement, 328–329
    - sources of ES cells and feeder cell lines, 325t
    - trypsin/EDTA, 330
  - epiblast stem cells (EpiSCs), 334
  - extra-embryonic endoderm stem (XEN) cells, 333–334
  - feeder cells for, 330–331
    - primary mouse embryonic fibroblasts (MEFs), 330
    - STO fibroblasts, 330–331
  - karyotype, 332–333
  - pathogen testing of, 331–332
  - progenitor cells from the CNS, 334
  - protocols, 338–404
    - chromosome counting, 363–365, 364f
    - cryopreservation of ES cells, 345–347
      - discussion, 347
      - freezing ES cells, 346
      - materials, 345
      - thawing ES cells, 346–347
    - derivation and culture of EpiSC cell lines, 382–388
      - cryopreservation, 386
      - culturing cells, 385–386
      - derivation, 383–385, 384f
      - materials, 382–383
      - method, 383–386
      - recipes, 386–388
    - derivation and culture of TS cell lines, 373–378
      - discussion, 376
      - materials, 373–374
      - method, 374–376, 375f
      - recipes, 376–378
    - derivation and culture of XEN cell lines, 379–381, 380f
    - derivation of ES cell lines using serum-containing culture conditions, 350–354, 351f–353f
    - derivation of ES cell lines using small-molecule inhibitors of Erk and Gsk3 signaling (2i), 355–362
      - embryo appearance, 358f
      - expansion of ES cell lines, 359, 359f
      - inner cell mass isolation by immunosurgery, 357–359
      - materials, 355–356
      - preimplantation epiblast preparation, 356–357
      - recipes, 360–362
      - summary diagram, 357f
    - differentiating ES cells into embryoid bodies by suspension culture, 368–369
    - differentiating ES cells into embryoid bodies in AggreWell plates, 370–372
    - feeder cell layer preparation from STO or MEF cells, 341–342
      - discussion, 342
      - materials, 341
      - mitomycin C treatment, 341–342
      - treatment by  $\gamma$ -irradiation, 342
    - isolation, culture, and differentiation of progenitor cells from central nervous system, 389–394
      - differentiation, 392
      - discussion, 392
      - dissection, 390–392, 391f
      - materials, 389–390
      - method, 390–392, 390f, 391f
      - recipes, 392–394
      - secondary neurospheres, 392
      - slide coating, 392
    - isolation, culture, and differentiation of radial glial-like neural progenitor cells in adherent cultures, 395–401
      - coating culture surfaces with AK-cyclo(RGDfC), 396
      - differentiation, 399, 400f
      - dissection, tissue dissociation, and cell seeding, 396–397
      - fixation and immunohistochemistry, 399
      - harvesting by trypsinization, 398–399
      - maintenance and propagation of cells, 397–398, 398f, 399f
      - materials, 395–396
      - methods, 396–399, 398f–400f
      - recipes, 400–401
    - passage of ES cells, 343–344
    - primary mouse embryonic fibroblasts (MEFs), preparation of, 338–340
      - discussion, 339–340
      - materials, 338–339
      - method, 339
      - testing serum batches, 348–349
    - trophoblast stem (TS) cells, 333–334
    - in vitro differentiation, 333
- Embryonic stem (ES) cell-mediated transgenesis, 440–442
- examples, 441–442
  - overview, 440–441
  - vectors (*see* Vector design)
  - in vitro screen for widespread transgene expression (protocol), 455–457
- Embryo recipients, 92–93
- establishing breeding colony to supply, 88–89
  - production of, 93
  - sterile male mice to induce pseudopregnancy, 93
- Embryo transfer
- overview, 199–203
    - background, 199–200
    - nonsurgical embryo transfer (NSET), 200

- number of embryos to transfer, 200–201  
 recipient females, 201–202  
 technical aspects, 202–203, 202f  
 unilateral and bilateral transfers, 201  
 zona pellucida role in, 200
- protocols  
 oviduct transfer of embryos, 211–215, 212f, 213f  
 uterine transfer of embryos, 216–219, 218f  
 for rederivation of animals into SPF colonies, 637–638  
 rederivation procedure for, 639
- Embryo transfer pipette  
 air bubbles in, 202–203, 202f  
 loading embryos into, 202, 203, 213, 217  
 reducing capillary action in, 202  
 structure of, 202
- EMMA (European Mouse Mutant Archive), 768
- ENCODE project, 1, 17
- Endoderm  
 anterior visceral, 52  
 formation of definitive, 45–49  
 gut, 68  
 parietal, 68–71, 69f–71f, 73, 73f  
 primitive, 38f, 41, 42f, 490  
 visceral, 68–71, 69f–71f, 73
- Endotoxin, as DNA prep contaminant, 248
- Enhanced cyan fluorescent protein (ECFP), 690f, 692
- Enhanced green fluorescent protein (EGFP), 244, 690f, 692
- Enhanced yellow fluorescent protein (EYFP), 692
- ENU (ethylnitrosourea), 16
- Enucleation  
 of zygotes, 579–580, 580f  
 pipette, 578–579, 579f, 585–586, 587f
- Environmental Protection Agency (EPA), U.S., 791
- Enzymes. *See also specific enzymes*  
 as DNA prep contaminant, 248  
 heat inactivation of, 248
- Enzyme solution  
 for adult tissue dissociation (recipe), 400  
 for neurosphere assay (recipe), 393
- Eosin, 744
- Ephrussi, Boris, 13
- Epiblast, 38f, 41–45, 43t, 44t  
 derivation of ES cell lines using small-molecule inhibitors of Erk and Gsk3 signaling (2i) (protocol), 355–362  
 fate maps of epiblast before and after gastrulation, 48f  
 gut development, 67–68  
 as pluripotent tissue, 44–45
- Epiblast stem cells (EpiSCs), 44, 334  
 derivation and culture of EpiSC cell lines (protocol), 382–388  
 cryopreservation, 386  
 culturing cells, 385–386  
 derivation, 383–385, 384f  
 materials, 382–383  
 method, 383–386  
 recipes, 386–388
- Epidermal growth factor (EGF), 389
- Epinephrine, 214
- EpiSC + FGF2 medium (recipe), 388
- EpiSC medium (recipe), 387
- EpiSCs. *See* Epiblast stem cells
- Equipment, 769–771  
 biological safety cabinets, 770–771  
 electrofusion apparatus, 776
- incubators, 769–770
- laminar flow hoods, 770
- for micromanipulation, 773–774  
 laser systems, 774  
 microinjection systems, 773–774  
 micromanipulators, 773  
 piezoelectric systems, 774
- microscopes, 771–773  
 inverted microscopes, 772  
 stereomicroscopes, 771  
 temperature controllers for heating and cooling stages, 772  
 vibration-free tables, 773
- microtool production from glass capillaries, 774–775  
 bevelers, 775  
 grinders, 775  
 microforges, 775  
 pullers, 775
- refrigerators and freezers, 769
- for spermatogonial stem cell injection into testis, 419–420, 419f
- surgical instruments, 776–777
- ES cells. *See* Embryonic stem (ES) cell(s); Embryonic stem (ES) cell lines
- Estrogen receptor, 239, 440
- Estrous cycle  
 indicators, 95t  
 induction by male pheromones (Whitten effect), 95  
 lighting cycle and, 94  
 phases of, 94, 95t, 104f  
 selecting mice in estrus, 103  
 stage determination by vaginal smear (protocol), 617–618, 618f  
 synchronization of, 95
- Estrus  
 indicators of, 95t  
 induction of, 95
- Ethanol  
 for parthenogenetic activation of oocytes, 572, 574–576, 575f  
 precipitation for contaminant removal in DNA prep, 248
- Ethidium bromide, as DNA prep contaminant, 248
- Ethylnitrosourea (ENU), 16
- European Conditional Mouse Mutagenesis Program (EuCOMM), 90
- European Mouse Mutant Archive (EMMA), 768
- Euthanasia, by cervical dislocation, 137, 137f
- Evans, Martin, 14, 18
- Exon, 432
- Expression. *See* Gene expression
- Extra-embryonic endoderm stem (XEN) cells, 333–334, 490  
 derivation and culture of XEN cell lines (protocol), 379–381, 380f
- Extra-embryonic membranes, isolating  
 overview, 150–151  
 protocol, 167–169, 168f
- Extra-embryonic tissues, 68–70, 69f, 70f  
 development stages, 26t–27t  
 differentiation of extra-embryonic mesoderm, 72, 72f
- EYFP (enhanced yellow fluorescent protein), 692
- F**
- Fate map  
 of early postimplantation embryo, 49  
 of epiblast before and after gastrulation, 48f
- FBS (fetal bovine serum), 328–329
- Feeder cells for ES cell lines, 325t, 330–331  
 preparation from STO or MEF cells (protocol), 341–342  
 discussion, 342  
 materials, 341  
 mitomycin C treatment, 341–342  
 treatment by  $\gamma$ -irradiation, 342
- primary mouse embryonic fibroblasts (MEFs), 330, 341–342  
 STO fibroblasts, 330–331, 341–342
- Feeder-conditioned medium (Feeder-CM) (recipe), 377
- Females  
 artificial insemination of, 597–598, 615–616, 616f, 619–622, 620f  
 dissection of reproductive organs, 138, 138f  
 embryo donors, 89–92  
 embryo recipients, 92–93  
 establishing breeding colony, 88–89  
 estrous cycle, 94–95, 95t, 103, 104t  
 housing of, 88  
 for pregnant females, 94  
 infertility, 594–595  
 isolating germ cells from postimplantation embryo genital ridge, 176–177, 177f  
 oogenesis, 31f  
 selecting mice in estrus and checking plugs (protocol), 103, 104f  
 sex determination and differentiation, 60–61  
 timing of germ-cell development, 51f  
 urogenital system, 594f
- Fertilization, 33f, 34–35  
 in vitro fertilization (IVF), 596–597, 604–614
- Fetal bovine serum (FBS), 328–329
- Fetal death, 151
- Fetal great vessel visualization by plastic coating, 690f, 694, 765–766
- Fetal vasculature visualization by India ink injection, 690f, 694, 763–764
- FGF (fibroblast growth factor)  
 FGF2 stock solution (10  $\mu$ g/mL) (recipe), 388  
 FGF4 stock solution (1000 $\times$ , 25  $\mu$ g/mL) (recipe), 377
- FHM medium, 111t, 128t, 130t
- FIAU, 465, 469t
- Fibroblasts  
 primary mouse embryonic fibroblasts (MEFs), 330  
 feeder cell layer preparation from (protocol), 341–342  
 preparation of (protocol), 338–340  
 discussion, 339–340  
 materials, 338–339  
 method, 339
- reprogramming  
 integrating *piggyBac* transposon transgenes into mouse fibroblasts by electroporation (protocol), 424–425  
 integrating *piggyBac* transposon transgenes into mouse fibroblasts using chemical methods (protocol), 426–427  
 reprogramming mouse fibroblasts with *piggyBac* transposons (protocol), 428–430, 429f
- Firefly luciferase gene, 244

- Fixation. *See* Tissue fixation
- Fixative for frozen sections (recipe), 749
- Fixative for staining intact embryos (recipe), 749
- Flp recombinase  
described, 439  
site-specific recombination-mediated insertions, 448–449
- Flunixin meglumine, 199, 199t
- Fluorescein isothiocyanate (FITC) filter sets, 693
- Fluorescent in situ hybridization (FISH)  
(FISH)-based chromosome painting, karyotyping by, 332, 554–555  
identification of homozygous transgenic mice/embryos, 556
- Fluorescent proteins  
protocols  
intracellular observation of fluorescent proteins in fixed cells, 756  
observation in living cells, 755  
visualization of fluorescent protein expression in whole-mount postimplantation-stage embryos, 753–754  
as reporters, 436–437, 437f  
for testing pluripotency and germline competence of induced pluripotent stem (iPS) cells, 408, 408f  
visualization techniques for cells using, 690f, 692–693  
compound, confocal, and deconvolution microscope use, 693  
dissecting microscope use, 692–693  
macroscopic observation, 692  
observation in living cells, 693, 755
- Fluorinert, 586
- Follicle-stimulating hormone (FSH), 30, 95
- Fostering, 94, 203, 220–221
- Foxa2* gene, 65
- Free radicals, 116
- Freezers, 769
- Freezing ES cells in 96-well plates (protocol), 475–477
- Freezing medium  
2× (recipe), 477  
for feeder cell culture (2×) (recipe), 388  
for feeder-free cell culture (1×) (recipe), 388
- FRT sites, 439
- FSH (follicle-stimulating hormone), 30, 95
- FuGENE HD transfection reagent, 426–427
- Furth, Jacob, 7
- Fused* (*Fu*) locus, 7
- Fyrite Kit, 328
- G**
- G418, 465, 465t, 469t, 472, 474
- γ-irradiation, feeder cell layer preparation from STO or MEF cells by, 342
- Ganciclovir, 469t
- Gardener, Richard, 13
- Gas containers  
for incubators, 770  
safe handling of, 790
- Gastrulation, fate maps of epiblast before and after, 48f
- G-banding, 332
- Gelase, 248
- Gelatin (0.1%) (recipe), 360
- Gene expression  
effect of embryo culture on, 117–118  
in vitro screen for widespread transgene expression (protocol), 455–457
- Gene knockout, 442–445
- Gene products, visualization techniques for, 684–691. *See also specific protocols*
- cutting sections, 685  
dehydrating in alcohol, 685  
dewaxing and rehydrating sections, 686  
embedding in wax, 685  
glass slide and coverslip preparation, 685  
immunohistochemistry  
of embryo sections, 686  
fluorescent analysis, 686  
general techniques for, 684–686  
of whole-mount embryos, 686, 689f  
isolating total RNA from mouse embryos or fetal tissues, 684
- in situ hybridization  
avoiding contamination with RNase, 687  
of embryo and tissue sections with RNA probes, 686–687  
general techniques for, 684–686  
of whole-mount embryos with RNA probes, 687–688, 689f, 691  
staining for alkaline phosphatase activity, 689f, 691  
staining for β-galactosidase activity, 689f, 691  
tissue fixation, 685
- Gene targeting  
described, 442–445  
detection and analysis of ES cell-mediated gene/genome alterations, 554
- Genetics, 1–11. *See also* Genome  
beginnings of mouse, 3  
of coat color, 75–77, 75t, 76f  
developmental  
heritage of experimental embryology, 11–14  
origin of, 9–11  
genome manipulation, 14–15  
systematic mutagenesis and phenotyping, 15–18
- Gene transfer. *See* Electroporation
- Gene trapping  
described, 17, 445–446  
detection and analysis of ES cell-mediated gene/genome alterations, 554
- Genital ridge, isolating germ cells from postimplantation embryos, 151–152, 152f, 175–177, 176f–177f
- Genome  
current annotation, 1  
linkage conservation and synteny with human genome, 2  
manipulation of, 14–15  
statistics, 4t
- Genomic and gene elements, in vector design, 432–434  
arms of homology, 433–434, 434f  
exon, 432  
internal ribosomal entry site (IRES), 432  
intron, 432  
polyadenylation signal, 432  
promoters, 432  
splice donor, 432  
2A peptide sequence, 433
- Genotyping, 551–570  
chimeras, 501, 501f, 502  
cloning transgene/host DNA junctions, 555  
detection and analysis of ES cell-mediated gene/genome alterations, 554–555  
detection and analysis of transgene from pronuclear injection, 552–553  
copy number, 553  
PCR genotyping, 552  
Southern blot analysis, 552–553  
ES cell colonies before picking (protocol), 485–487  
identification of homozygous transgenic mice/embryos, 555–557  
PCR analysis using flanking primer, 557f  
in situ hybridization to interphase nuclei, 556  
Southern blot analysis using flanking probe, 556  
test breeding, 556
- protocols  
DNA preparation from cultured cells for Southern blot or PCR, 567–568  
isolation of high-molecular weight DNA from embryonic or extra-embryonic tissues, 560–561  
isolation of high-molecular weight DNA from mouse tail tips, 558–559  
PCR template preparation from embryonic and adult tissue samples, 564–565  
PCR template preparation from mouse tail tissue, 566  
polymerase chain reaction (PCR), 567–568  
tissue lysate preparation for PCR template, 562–563  
tissue samples for, 551–552
- Genotyping assays, 462, 462t
- Germ cell isolation from genital ridge of postimplantation embryos  
overview, 151–152, 152f  
protocol, 175–177, 176f–177f
- Germ layer explant recombinant culture  
overview, 151  
protocol, 173–174
- Germ layers, separating postimplantation  
overview, 151  
protocol, 170–172, 171f
- Germline, origin of, 49–51, 50f
- Germline-competent stem cells derived from adult mice, 403–430  
overview, 403–409  
induced pluripotent stem (iPS) cells, 406–408  
reprogramming vectors, 407–408, 407f  
testing pluripotency and germline competence of, 408, 408f  
spermatogonial stem cells (SSCs), 404–406  
culture of, 404–405  
transplantation of, 405–406, 405f
- protocols, 410–430  
cryopreserving and thawing spermatogonial stem cells, 416–417  
culture and expansion of primary undifferentiated spermatogonial stem cells, 413–415



- establishment of primary culture, 413–414, 415f  
 expansion of primary undifferentiated culture, 414–415  
 materials, 413  
 integrating *piggyBac* transposon transgenes into mouse fibroblasts by electroporation, 424–425  
 integrating *piggyBac* transposon transgenes into mouse fibroblasts using chemical methods, 426–427  
 isolation of the spermatogonial stem cell-containing fraction from testes, 410–412  
 materials, 410–411  
 percoll isolation, 412  
 testes disaggregation, 411  
 $\text{Thy1}^+$  selection by magnetic-activated cell sorting, 412  
 reprogramming mouse fibroblasts with *piggyBac* transposons, 428–430, 429f  
 spermatogonial stem cell transplantation to the testis, 418–423  
 busulfan treatment of adult recipients, 419  
 discussion, 422  
 injection of cells into recipient tissue, 420–422, 421f  
 materials, 418–419, 419f  
 preparation of cells for transplantation, 420  
 preparation of injection pipette and injector, 419–420  
 Gestation  
   development staging, 24–27, 25f, 26t–27t, 28f–30f  
   length of, 24  
 GFP. *See* Green fluorescent protein  
 Giemsa staining of chromosomes, 332  
 Gittschier buffer (recipe), 486  
 Glasgow Minimal Essential Medium (G-MEM), 327  
 Glass capillaries, microtool production from, 774–775  
 Glass slide and coverslip preparation for in situ hybridization, 685, 704–705  
 Glucose, in embryo culture media, 113–114  
 Gluecksohn-Waelsch, Salome, 10  
 Glutamine  
   L-alanyl-L-glutamine, 114  
   sperm cryopreservation using cryoprotectant containing L-glutamine (protocol), 671–675, 672f–674f  
 G1 medium, 111t, 112  
 G2 medium, 111t, 112  
 Gonadotropin  
   dose, 96t, 97  
   preparation of, 105  
   time of administration, 97  
 Gonads. *See also* Ovary; Testis  
   sex determination and differentiation, 60–61  
 Gordon, Jon, 14  
 Gorer, Peter, 7  
 Gossler, Achim, 15  
 Graafian follicles, 30, 32f  
 Granulosa cells, luteinized, 32  
 Great vessel visualization by plastic coating, 690f, 694, 765–766  
 Green fluorescent protein (GFP), 692  
   fixation and paraffin embedding for GFP visualization (protocol), 757–758  
   intracellular observation in fixed cells (protocol), 756  
   observation in living cells (protocol), 755  
   as reporter gene in transgene constructs, 244, 437  
 Grinders, 775  
 Gropp, Alfred, 9  
 Growth factors, in embryo culture media, 117  
 Grüneberg, Hans, 10  
 Gurdon, John, 18, 403, 406  
 Gut development, 67–68  
 Gynogenotes, 571  
  
**H**  
 Hadorn, Ernst, 10, 12  
 Hair follicle, 74–75, 74f  
 Haldane, J.B.S., 3, 10  
 Hammond, John, Jr., 12  
 Handling blastocysts for fixation (protocol), 700  
 Hazardous chemicals, properties of common, 793–794  
 HB-EGF (heparin-binding epidermal growth factor), 117  
 hCG (human chorionic gonadotropin), 91, 95, 97, 105, 315  
 Heavy inhibitory solution (HI) (recipe), 393  
*Helicobacter typhlonius*, 639  
 Hensen's node. *See* Node  
 HEPA (high-efficiency particulate air)-filtration, 87, 770  
 Heparin-binding epidermal growth factor (HB-EGF), 117  
 Heparin stock solution (1000 $\times$ , 1 mg/mL) (recipe), 377  
 Hepatectomy, partial, 227–228  
 HEPES-buffered DMEM with FBS (recipe), 166, 169, 172  
 HEPES-buffered media for embryo handling, 128–129, 128t  
 Hermaphrodites, 500  
 Herrmann, Bernhard, 11  
*Hex* gene, 45  
 High-efficiency particulate air (HEPA)-filtration, 87, 770  
 High-osmolality vitrification, embryo cryopreservation by, 658–664, 660f, 661f  
   cooling embryos, 659–660, 662  
   discussion, 661  
   in insemination straws, 662–664  
   loading embryos into cryotube, 660f  
   materials, 658–659  
   warming embryos, 660–661, 661f, 662–663  
 Hind limb bud morphology, 152f  
 Histocompatibility genes, 8  
 Histone H4 promoter, 245  
 HMGR promoter, 245  
 Holding pipette  
   for ES cell chimera formation, 506  
   for intracytoplasmic sperm injection (ICSI), 624  
   making holding pipettes (protocol), 294–295  
   in microinjection setup, 299–300, 300f  
   pressure control in, 259  
   for pronuclear microinjection, 256–257, 256f, 257f, 294–295  
 Homologous recombination  
   arms of homology and, 433–434, 434f  
   during pronuclear microinjection, 241  
   replacement by, 446–448  
     hit and run example, 447  
     *piggyBac* transposon example, 448  
     tag and exchange example, 447  
 Homology search, 15–16  
 Homozygous  
   ES cells for mutations, 451–452  
   transgenic mice/embryos, identifying, 555–557  
 Honolulu method of SCNT, 572  
 Housekeeping genes, promoters of, 245  
 Housing  
   females for embryo transfer, 88  
   males, 88  
   pregnant females, 94  
*Hox* genes, 58, 59f, 63, 64f  
 hPAP (human placental alkaline phosphatase) gene, 244, 437  
*hprt* (hypoxanthine phosphoribosyltransferase), 15, 435, 436  
 HSV-tk (herpes simplex virus-1 thymidine kinase gene), 436, 465  
 HTF (human tubal fluid), 597  
 Human chorionic gonadotropin (hCG), 91, 95, 97, 105, 315  
 Human Genome Project, 2  
 Human menopausal gonadotropin, 95  
 Human placental alkaline phosphatase (hPAP) gene, 244, 437  
 Human tubal fluid (HTF), 597  
 Human tubal fluid medium (recipe), 612, 622  
 Human tubal fluid medium with calcium (recipe), 613  
 Hyaluronidase  
   cumulus cell removal with, 34, 141  
   release from acrosome, 34  
 Hyaluronidase (recipe), 781  
 Hybridization mix (recipe), 739  
*hygro* (hygromycin-resistance gene), 435, 465t  
 Hygromycin B, 435, 465t, 469t  
  
**I**  
 Ice crystals, 634  
 ICM. *See* Inner cell mass  
 ICR mice, 590  
 ICSI. *See* Intracytoplasmic sperm injection  
 Identification. *See* Animal identification  
 IETS (International Embryo Transfer Society), 638  
 IGF-II (insulin-like growth factor II), 117  
 IGTC (International Gene Trap Consortium), 768  
 IKMC (International Knockout Mouse Consortium), 17, 90, 492, 768  
 Imaging  
   embryos after whole-mount in situ hybridization, 741  
   mounting embryos for microscopic observation and imaging (protocol), 726, 727f  
   static culture of postimplantation embryos for, 154, 184–186, 185f  
 IMDM (Iscove's Modified Dulbecco's Medium), 327  
 Immunofluorescent staining of whole-mount embryos (protocol), 724–725  
 Immunohistochemistry  
   overview  
   of embryo sections, 686

- Immunohistochemistry (*Continued*)  
 fluorescent analysis, 686  
 general techniques for, 684–686  
 of whole-mount embryos, 686, 689f  
 protocols  
 of embryo sections, 715–718  
 of whole-mount embryos, 719–721
- Immunosurgery, isolating inner cell mass by, 357–359, 358f, 549–550, 549f
- IMPC (International Mouse Phenotyping Consortium), 1, 768
- Implantation  
 events in, 37, 39, 40f  
 timing of, 149, 150  
 visualizing sites, 150, 150f, 157–158
- IMSR (International Mouse Strain Resource), 768
- Incisions, surgical, 196
- Incubators, 116, 769–770
- India ink injection, vasculature visualization by, 690f, 694, 763–764
- Induced pluripotent stem (iPS) cells  
 overview, 14, 406–408  
 reprogramming vectors, 407–408, 407f  
 testing pluripotency and germline competence of, 408, 408f  
 protocols  
 integrating *piggyBac* transposon transgenes into mouse fibroblasts by electroporation, 424–425  
 integrating *piggyBac* transposon transgenes into mouse fibroblasts using chemical methods, 426–427  
 reprogramming mouse fibroblasts with *piggyBac* transposons, 428–430, 429f  
 Yamanaka factors, 406, 407
- Inducible systems, 439–440  
 tamoxifen, 440  
 tetracycline, 440, 458–459  
 in vitro screen to identify silent but activatable integration sites for tet-inducible transgene (protocol), 458–460
- Inhalation anesthetics, 198
- Inhibitors (2i), 10 mM stock (recipe), 360
- Inhibitors of Erk and Gsk3 signaling (2i)  
 derivation of ES cell lines using (protocol), 355–362  
 embryo appearance, 358f  
 expansion of ES cell lines, 359, 359f  
 inner cell mass isolation by immunosurgery, 357–359  
 materials, 355–356  
 preimplantation epiblast preparation, 356–357  
 recipes, 360–362  
 summary diagram, 357f
- Injection buffer preparation  
 for BAC/YAC DNA (protocol), 290  
 for standard-sized DNA (protocol), 289
- Injection chambers, for pronuclear microinjection, 260–262, 260f–262f, 299, 299f
- Injection pipette  
 in cloning mice procedure, 585–586, 587f  
 for ES cell chimera formation, 506  
 for intracytoplasmic sperm injection (ICSI), 624  
 making injection pipettes (protocol), 296–297  
 positioning of, 299, 300f
- pressure control in, 259–260  
 for pronuclear microinjection, 257–258, 296–297  
 for pronuclear transfer, 578–579, 579f
- Inner cell mass (ICM), 36–37, 37f  
 disaggregating cleavage-stage embryos and inner cell mass of blastocysts into individual cells (protocol), 547–548  
 injection into blastocyst to create chimeras, 490  
 isolating inner cell mass of blastocysts (protocol), 549–550, 549f  
 isolation by immunosurgery, 357–359, 358f, 549–550, 549f
- Insertional mutagenesis, 17
- Insertions, site-specific recombination-mediated, 448–449
- In situ hybridization  
 fluorescent in situ hybridization (FISH) (FISH)-based chromosome painting, karyotyping by, 332, 554–555  
 identification of homozygous transgenic mice/embryos, 556  
 overview  
 avoiding contamination with RNase, 687  
 of embryo and tissue sections with RNA probes, 686–687  
 general techniques for, 684–686  
 of whole-mount embryos with RNA probes, 687–688, 689f, 691  
 protocols  
 dewaxing and rehydrating sections before in situ hybridization or staining, 714  
 of embryo and tissue sections with radiolabeled RNA probes, 728–732  
 autoradiography, 733–734  
 hybridization, 731  
 materials, 728–729  
 posthybridization washing, 731–732  
 prehybridization, 730–731  
 probe preparation, 729–730  
 glass slide and coverslip preparation, 704–705  
 imaging embryos after whole-mount in situ hybridization, 741  
 sectioning embryos after whole-mount in situ hybridization, 742–743  
 of whole-mount embryos with nonradiolabeled RNA probes, 735–740  
 discussion, 738–739  
 embryo preparation, 736  
 histochemistry, 738  
 materials, 735–736  
 post-antibody washes, 738  
 posthybridization washes, 738  
 pretreatment and hybridization, 737  
 probe preparation, 736–737  
 recipes, 739–740
- Institutional safety office, 789
- Insulin-like growth factor I (IGF-I), 117
- Insulin-like growth factor II (IGF-II), 117
- Internal ribosomal entry site (IRES), 243, 408, 432
- International Embryo Transfer Society (IETS), 638
- International Gene Trap Consortium (IGTC), 768
- International Knockout Mouse Consortium (IKMC), 17, 90, 492, 768
- International Mouse Phenotyping Consortium (IMPC), 1, 768
- International Mouse Strain Resource (IMSR), 768
- Internet resources, 785–787
- Intestinal tract, development stages, 26t–27t
- Intrabursal transfer of sperm (ITS), 620
- Intracytoplasmic sperm injection (ICSI), 598, 623–630, 626f, 627f  
 overview, 598  
 protocol, 623–630, 626f, 627f
- Intraperitoneal injection (protocol), 107–108, 107f
- Intrauterine growth restriction (IUGR), 151
- Introns, 245, 432
- Inverse PCR, 555
- Inversions, induced site-specific, 450
- Inverted microscopes, 772
- In vitro differentiation of ES cell lines, 333
- In vitro fertilization (IVF), 596–597, 604–614  
 discussion, 611–612  
 Jackson Laboratory method, 605–608  
 laser assisted, 597  
 materials, 604–605  
 media for, 596–597  
 Nakagata method, 608–610  
 overview, 596–597  
 protocol, 604–614  
 recipes, 612–614  
 troubleshooting, 611
- In vitro screen to identify silent but activatable integration sites for tet-inducible transgene (protocol), 458–460
- In vitro–translated RNA for microinjection, preparation of (protocol), 292–293
- Ion-exchange chromatography, for contaminant removal in DNA prep, 248
- IPS cells. *See* Induced pluripotent stem (iPS) cells
- IRES (internal ribosomal entry site), 243, 408, 432
- Iscove's Modified Dulbecco's Medium (IMDM), 327
- Isoflurane, 198
- ITS (intrabursal transfer of sperm), 620
- IUGR (intrauterine growth restriction), 151
- IVF. *See* In vitro fertilization
- ## J
- Jackson Laboratory, 7, 8, 12, 13, 493, 768
- Jacob, François, 13
- Jaenisch, Rudolf, 14
- ## K
- Karyoplast, 572, 579, 580f, 581–582, 581f
- Karyotype/karyotyping  
 chromosomal banding, 332  
 chromosome counting (protocol), 363–365, 364f  
 embryonic stem (ES) cell lines, 332–333  
 fluorescent in situ hybridization (FISH)-based chromosome painting, 332, 554–555  
 spectral karyotyping (SKY), 332
- Kaufmann, Matt, 14

- Ketamine/xylazine, 198, 198t  
Ketoprofen, 199, 199t  
Kidney  
  development of, 60  
  nephrectomy, 228–229  
  tissue transplantation under capsule of (protocol), 222–224, 223f  
Klenow fragment of DNA polymerase I, 739  
KLF4 transcription factor and gene, 406, 407  
KnockOut D-MEM medium, 327  
KnockOut Mouse Project (KOMP), 90  
KnockOut Serum Replacement (KSR), 324, 354  
KSOM + 2i inhibitors (100× stock) (recipe), 360  
KSOM medium, 111t, 112, 113, 127t, 130t, 534, 576
- L**  
Laboratory location, 87–88  
*lacZ* gene, 244  
  as reporter, 436, 440  
  for testing pluripotency and germline competence of induced pluripotent stem (iPS) cells, 408, 408f  
L-alanyl-L-glutamine, 114  
Laminar flow hoods, 770  
Laser safety, 790  
Laser systems, 774  
Lathrop, Abbie E.C., 4  
Lentivirus  
  infecting embryos by coculturing with lentivirus (protocol), 312–314  
  cleanup, 313  
  discussion, 314  
  embryo preparation, 312–313  
  materials, 312  
  setup for infection, 313  
  troubleshooting, 313  
  microinjection of lentivirus into embryos (protocol), 308–311  
  cleanup, 310  
  discussion, 311  
  injection of embryos, 309–310  
  materials, 308–309  
  setup, 309  
  troubleshooting, 310–311  
  vectors for pronuclear microinjection, 241, 308  
Lesch-Nyhan disease, 15  
Leukemia-inhibitory factor (LIF), 39, 117, 329–330  
L-glutamine, 114  
  sperm cryopreservation using cryoprotectant containing L-glutamine (protocol), 671–675, 672f–674f  
LH (luteinizing hormone), 30, 95, 97  
LIF (leukemia-inhibitory factor), 39, 117, 329–330  
Light exposure, embryo culture and, 117  
Light inhibitory solution (recipe), 394  
Light sources, 693, 771  
*limb deformity*, 11  
Limb formation, 61  
Lin, Teh Ping, 14  
Linkage, 3  
Little, Clarence C., 4, 7, 8  
Location of embryo manipulation laboratory, 87–88  
Loeb, Leo, 4, 7  
*loxP* sites, 239, 437–438, 444–445. *See also* Cre recombinase  
Luteinizing hormone (LH), 30, 95, 97  
Lyon, Mary, 10  
Lysis buffer (recipe), 484, 565, 570
- M**  
MABT (recipe), 739  
Males  
  establishing breeding colony, 88–89  
  estrus induction by male pheromones, 95  
  fighting among, 88  
  housing of, 88  
  infertility, 595–596  
  isolating germ cells from postimplantation embryo genital ridge, 176–177, 177f  
  sex determination and differentiation, 60–61  
  spermatogenesis, 31f, 32–34  
  sperm count depression after mating, 91  
  sterile, to induce pseudopregnancy, 93  
  studs for embryo donors, 91  
  timing of germ-cell development, 51f  
  urogenital system, 594f  
Mannitol, 539–541, 540f  
Mannitol (0.3 M) (recipe), 781  
MAP (mouse antibody production) testing, 331  
Markers, selectable, 434–436  
Martin, Gail, 14  
Material safety data sheets (MSDSs), 789  
Maternal leukemia inhibitory factor (LIF), 117  
Matings  
  to induce pseudopregnancy, 93  
  natural, 91  
  estrus indicators, 95t  
  lighting for, 95t  
  setting up, 94–95  
  stud mice, 91  
Matrigel Basement Membrane, 203, 225  
MBCD (methyl- $\beta$ -cyclodextrin), 597, 607, 609, 671  
McLaren, Anne, 12  
MCMV (mouse cytomegalovirus), 639  
Media. *See* Culture media; *specific media*  
Medical Pathological Waste (MPW), 791  
MEFs. *See* Primary mouse embryonic fibroblasts (MEFs)  
Melanoblasts, 75  
Meloxicam, 199, 199t  
Mendelian inheritance, 3  
Mercury, 586, 587f  
Mesoderm, 43, 43t  
  differentiation of extra-embryonic, 72, 72f  
  formation of, 45–49  
  generation of regional diversity in, 51–52  
  intermediate, 58, 60  
  of placenta, 73  
Mesonephros, 176f  
Metallothionein promoter, 245  
Metestrus 1, 95t  
Metestrus 2, 95t  
Methyl- $\beta$ -cyclodextrin (MBCD), 597, 607, 609, 671  
Methylene blue solution (recipe), 781  
MgCl<sub>2</sub> (1 M) (recipe), 394  
MGI (Mouse Genome Informatics) database, 4  
MHV (mouse hepatitis virus), 86, 331, 638  
Microchip implants, 100  
Microdrop culture method, 110, 119  
  setting up (protocol), 132–133, 133f  
Microforges, 775  
Microinjection. *See also* Pronuclear microinjection; *specific protocols*  
  of lentivirus into embryos (protocol), 308–311  
  preparation of DNA for, 252–254  
  DNA concentration determination, 253–254  
  filtration of DNA solution, 254  
  quality issues, 252–253  
  storage of DNA preparation, 254  
  of spermatogonial stem cells into testis, 420–422, 421f  
  of zygotes  
  overview, 262–263  
  protocol, 301–305, 303f  
  discussion, 304–305  
  materials, 301  
  method, 301–304, 303f  
  with RNA (protocol), 306–307  
Microinjection setup (protocol), 298–300, 299f, 300f  
Microinjection systems, 773–774  
Micromanipulation laboratory, setting up, 767–778  
  cryopreservation, 777  
  equipment, 769–771  
  biological safety cabinets, 770–771  
  electrofusion apparatus, 776  
  incubators, 769–770  
  laminar flow hoods, 770  
  for micromanipulation, 773–774  
  laser systems, 774  
  microinjection systems, 773–774  
  micromanipulators, 773  
  piezoelectric systems, 774  
  microscopes, 771–773  
  inverted microscopes, 772  
  stereomicroscopes, 771  
  temperature controllers for heating and cooling stages, 772  
  vibration-free tables, 773  
  microtool production from glass capillaries, 774–775  
  bevelers, 775  
  grinders, 775  
  microforges, 775  
  pullers, 775  
  refrigerators and freezers, 769  
  surgical instruments, 776–777  
  general considerations, 767–768  
  mouse strains for, 768  
Micromanipulators, 259, 298–299, 585f, 586–587, 586f, 773  
Microscopes, 771–773  
  inverted microscopes, 772  
  for pronuclear microinjection, 258  
  stereomicroscopes, 771  
  temperature controllers for heating and cooling stages, 772  
  vibration-free tables, 773  
Microtome  
  cutting wax sections using (protocol), 706–707  
  transferring sections onto water droplets (protocol), 708  
  vibratome, cutting thick sections using (protocol), 709–711  
  fixed specimens, 710–711  
  live specimens, 712–713  
  materials, 709, 712

- Microtool production from glass capillaries, 774–775  
 bevelers, 775  
 grinders, 775  
 microforges, 775  
 pullers, 775
- Microwave safety, 790
- Mid-gastrula organizer (MGO), 54
- Min* (multiple intestinal neoplasia), 16
- Miniroller apparatuses, 770
- Mintz, Beatrice, 12–13, 14, 490
- Mitomycin C treatment, for feeder cell layer preparation from STO or MEF cells, 341–342
- Mitotic arrestants, 332
- M2 medium, 110, 111t, 130t, 537–538, 537f, 540–541, 575
- M16 medium, 110, 111t, 126, 127t, 130t, 575, 576
- MMESSAGE mMACHINE kit, 292–293
- MMRRC (Mouse Mutant Regional Resource Centres), 768
- MMV (mouse minute virus), 331, 639
- Modified PBS supplemented with glucose, pyruvic acid, BSA, and antibiotics (recipe), 613
- Modular incubator chambers, 770
- Moloney murine leukemia virus (Mo-MLV), 14
- Monothioglycerol (MTG), sperm cryopreservation using cryoprotectant containing, 665–670
- Montoliu, Lluís, 493
- Morgan, Thomas Hunt, 3, 10
- Morphological mutants, 78, 78f
- Morula  
 aggregation, 543  
 compaction, 35–36  
 cryopreservation of embryos, 632, 652  
 eight-cell uncompact, 35  
 ES cell aggregation with to create chimeras, 490–493
- Morula-stage embryo collection (protocol), 143–145, 144f–145f
- Mounting embryos for microscopic observation and imaging (protocol), 726, 727f
- Mouse antibody production (MAP) testing, 331
- Mouse cytomegalovirus (MCMV), 639
- Mouse embryonic fibroblasts. *See* Primary mouse embryonic fibroblasts (MEFs)
- Mouse Genome Informatics (MGI) database, 4
- Mouse Genomes Project, 1
- Mouse hepatitis virus (MHV), 86, 331, 638
- Mouse HMGCR promoter, 245
- Mouse lines. establishment and maintenance of, 263–264
- Mouse metallothionein promoter, 245
- Mouse minute virus (MMV), 331, 639
- Mouse Mutant Regional Resource Centres (MMRRC), 768
- Mouse parvovirus (MPV), 638
- Mouse strains. *See* Strains
- MPF (M-phase-promoting factor), 572
- M-phase-promoting factor (MPF), 572
- MPV (mouse parvovirus), 638
- MPW (Medical Pathological Waste), 791
- MSDSs (material safety data sheets), 789
- MTF (mouse tubal fluid) medium, 111t, 112–113
- MTG (monothioglycerol), sperm cryopreservation using cryoprotectant containing, 665–670
- Müllerian duct, 60–61
- Mutagenesis  
 ethylnitrosourea (ENU), 16  
 gene-trap, 462  
 insertional, 17  
 systematic mutagenesis and phenotyping, 15–18
- Mycoplasma*  
 contamination of cell cultures by, 331  
 detection of, 331–332  
*M. pulmonis*, 638
- Myo5a* gene, 9
- Myotome, 57
- N**
- NaCl/ethanol mixture (recipe), 482
- NANOG, for testing pluripotency and germline competence of induced pluripotent stem (iPS) cells, 408, 408f
- National Institute of Environmental Health and Human Services, 793
- Natural matings, 91  
 estrus indicators, 95t  
 lighting for, 95t  
 setting up, 94–95, 95t
- N2B27 medium (recipe), 361
- N2B27 medium with LIF and 2i inhibitors (recipe), 361
- n*-butanol extraction, for contaminant removal in DNA prep, 248
- NDS buffer (100%) (recipe), 276–277
- Needles, sharpening tungsten by electrolysis, 160f
- Negative selectable markers, 436
- neo* (neomycin phosphotransferase gene), 15, 434–435, 465, 465t
- Neon electroporation system, 424, 463
- NEP basal medium (recipe), 394
- NEP (neuroepithelial) cells, isolation, culture, and differentiation of (protocol), 389–394
- NEP complete medium (recipe), 394
- Nephrectomy, 228–229
- Nervous system  
 development of, 26t–27t, 61–62  
 isolation, culture, and differentiation of progenitor cells from central nervous system (protocol), 389–394
- Neural folds, 56f, 57f, 62
- Neural plate, 61–62
- Neural tube development, 57f, 62–65  
 regional diversification, 62–65, 63f, 64f
- Neurobasal + B27 medium (recipe), 360
- Neurobasal medium + PSG (recipe), 394
- Neuroepithelial (NEP) cells, isolation, culture, and differentiation of (protocol), 389–394
- Neuronal cells, differentiation of RGLs into, 399
- Neurosphere assay (NSA), 322, 334, 389
- Neurospheres  
 formation of, 391f  
 secondary, 392
- Neurulation, 61–62
- Neutralization buffer (recipe), 734
- Nieuwkoop, Pieter, 12
- nodal*, 17
- Node, 52–54, 53f
- Nonobese diabetic (NOD) mouse strain, 324, 325t, 355, 493
- Nonsteroidal antiinflammatory drugs (NSAIDs), 199, 199t
- Nonsurgical embryo transfer (NSET), 200
- North American Conditional Mouse Mutagenesis Program (NorCOMM), 90
- Notch signaling pathway, 55
- NSA (neurosphere assay), 322, 334, 389
- NSAIDs (nonsteroidal antiinflammatory drugs), 199, 199t
- NSET (nonsurgical embryo transfer), 200
- N2 supplement  
 solution (100×) (recipe), 361  
 (100×) stock solutions (recipe), 361
- NTMT (recipe), 739, 780
- NucleoBond  
 isolating BAC DNA from bacterial cultures (protocol), 271–272  
 kits, 250t
- O**
- Occupational Safety and Health Administration (OSHA), 789
- OCT4 transcription factor and gene, 406, 407
- Oligodendrocyte cells, differentiation of RGLs into, 399
- 129 substrains, 491–492, 494f–496f, 494t–496t, 507–508
- Oocyte  
 collection for in vitro fertilization, 607, 609–610  
 cryopreservation, 636  
 fertilization, 33f, 34–35  
 maturation, 27, 30, 31f–32f  
 ovulation, 30, 32, 33f, 34t  
 parthenogenetic activation of oocytes, 572, 574–576, 575f, 576f  
 partial zona dissection (PZD) of, 597  
 properties of ovulated, unfertilized, 34t
- Oogenesis, 31f
- Origin of the laboratory mouse, 4, 5f–7f
- OSHA (Occupational Safety and Health Administration), 789
- Osmolarity, of embryo culture media, 115
- Ovariectomy  
 overview, 203–204  
 protocol, 232–233  
 materials, 232  
 method, 232–233
- Ovary  
 cryopreservation, 637, 676–678  
 removal (*see* Ovariectomy)  
 transplantation, 596, 601–603, 602f  
 overview, 596  
 protocol, 601–603, 602f
- Oviduct  
 dissection for zygote removal, 139–142, 140f–141f  
 two-cell- to morula-stage embryo collection from, 143–145, 144f–145f
- Oviduct transfer of embryos (protocol), 211–215, 212f, 213f  
 discussion, 215  
 materials, 211  
 method, 211–214, 212f, 213f
- Ovulation, 30, 32  
 signs of, 201
- Oxygen tension, for embryo culture, 115–116

**P**

- PAF (platelet activating factor), 117  
Pancreatin/trypsin solution for separating germ and tissue layers (recipe), 781  
Paracetamol, 199  
Paraffin oil, 122, 202  
Paraffin wax  
  embedding for GFP visualization (protocol), 757–758  
  types of, 685  
Paraformaldehyde 4%, preparation of (protocol), 698  
Paraplast, 685  
Parietal endoderm, 68–71, 69f–71f, 73, 73f  
Parietal yolk sac  
  development, 68–69, 69f  
  isolation of, 151  
Partial zona dissection (PZD) of oocytes, 597  
Passage of ES cells (protocol), 343–344  
Passaging and freezing ES cells in 96-well plates (protocol), 475–477  
Pathogenesis  
  overview, 571–572  
  protocol, 574–576, 575f, 576f  
Pathogen control in mouse colony, 86–87  
Pathogen testing of embryonic stem (ES) cell lines, 331–332  
PBS (phosphate-buffered saline). *See* Phosphate-buffered saline  
PBSMT for immunohistochemistry of whole-mount embryos (recipe), 782  
PBT (recipe), 782  
PCR  
  buffer (1×) (recipe), 486  
  buffer (10×) (recipe), 568  
  genotyping  
    cloning transgene/host DNA junctions, 555  
    for detection and analysis of ES cell-mediated gene/genome alterations, 554  
    for detection and analysis of transgene from pronuclear injection, 552  
  DNA preparation protocols for  
    isolation of high-molecular weight DNA from embryonic or extra-embryonic tissues, 560–561  
    isolation of high-molecular weight DNA from mouse tail tips, 558–559  
    preparation from cultured cells, 567–568  
  ES cell colonies before picking (protocol), 485–487  
  identification of homozygous transgenic mice/embryos using flanking primer, 557, 557f  
  protocol for polymerase chain reaction (PCR), 567–568  
  tissue lysate preparation protocols for PCR templates  
    from embryonic, yolk sac, and adult tissues, 562–563  
    from embryonic and adult tissue samples, 564–565  
    from mouse tail tissue, 566  
  lysis buffer (recipe), 486  
  pathogen testing of embryonic stem (ES) cell lines, 331–332  
  reaction mixture (recipe), 487  
  Splinkerette, 555  
  tissue homogenization buffer (10×) (recipe), 563  
Percoll, 412  
PFGE. *See* Pulsed-field gel electrophoresis  
PGCs. *See* Primordial germ cells  
pH, of embryo culture media, 116  
Pharyngeal region, development of, 66–67, 67f  
Phenol, as DNA prep contaminant, 248  
Phenol:chloroform extraction, for contaminant removal in DNA prep, 248  
Phenol:chloroform solution (recipe), 782  
φC31 integrase  
  described, 439, 463  
  site-specific recombination-mediated insertions, 448–449  
Phosphate-buffered saline  
  recipe, 782–783  
  at pH 7.4 with Ca<sup>2+</sup>/Mg<sup>2+</sup> (recipe), 740  
  at pH 7.4, 10× (recipe), 740  
Pierce, Barry, 13  
Piezoelectric systems, 774  
piggyBac transposon, 17, 241, 243, 407, 448  
  integrating piggyBac transposon transgenes into mouse fibroblasts by electroporation (protocol), 424–425  
  integrating piggyBac transposon transgenes into mouse fibroblasts using chemical methods (protocol), 426–427  
  reprogramming mouse fibroblasts with piggyBac transposons (protocol), 428–430, 429f  
Pink-eyed dilution, 3, 76f, 77, 507  
Pipette  
  braking, 222, 223, 223f  
  in cloning mice procedure, 585–586, 587f  
  for embryo-handling, 121–122, 122f  
  components of, 121–122, 122f  
  hand-controlled, 121–122, 122f  
  making from hard glass capillaries, 134–135  
  reducing capillary action in, 122  
  siliconized, 136  
  embryo transfer  
    air bubbles in, 202–203, 202f  
    loading embryos into, 202, 203, 213, 217  
    reducing capillary action in, 202  
    structure of, 202  
  enucleation pipette, 578–579, 579f, 585–586, 587f  
  from glass capillaries, 134–135, 774–775  
  holding pipette  
    for ES cell chimera formation, 506  
    for intracytoplasmic sperm injection (ICSI), 624  
    for pronuclear microinjection making holding pipettes (protocol), 294–295  
    in microinjection setup, 299–300, 300f  
    pressure control in, 259  
    for pronuclear microinjection, 256–257, 256f, 257f, 294–295  
  injection pipette  
    in cloning mice procedure, 585–586, 587f  
    for ES cell chimera formation, 506  
    for intracytoplasmic sperm injection (ICSI), 624  
  making ES cell injection needles (protocol), 514–517, 514f, 516f  
  for pronuclear microinjection making injection pipettes (protocol), 296–297  
  positioning of, 299, 300f  
  pressure control in, 259–260  
  for pronuclear microinjection, 257–258, 296–297  
  for pronuclear transfer, 578–579, 579f  
  siliconized, preparing, 136  
  for surgical artificial insemination, 620f  
Placenta, 68, 69f, 72–74, 73f  
Plasmid rescue, 555  
Plastic coating, great vessel visualization by, 690f, 694, 765–766  
Platelet activating factor (PAF), 117  
Plugging rate, 315  
Plugs, 103, 104f  
Pluripotent cells. *See* Induced pluripotent stem (iPS) cells  
PMSG (pregnant mare serum gonadotropin), 91, 95, 97, 105–106, 315  
Polyadenylation signal, 432  
Polyamine mix (1000×) (recipe), 284  
Polyamines, 274, 290–291  
Poly-L-lysine, 685, 705  
Polymerase chain reaction (PCR). *See* PCR  
Polyspermy, 34  
Polyvinylpyrrolidone (10%–12%) (recipe), 590  
Polyvinylpyrrolidone, use in cloning mice protocol, 586–588, 587f  
Pontamine Sky Blue, 157  
Positive selectable markers, 434–435  
Postimplantation embryos, 149–193  
  culture of  
    overview, 152–154  
    preparing embryos, 152–153  
    roller culture, 153, 178–181, 179f–180f  
    static culture, 153–154, 182–186  
    for imaging, 154, 184–186, 185f  
  development of (*see* Development)  
  dissection, 771  
  electroporation  
    overview, 154–155, 155f  
    in utero (protocol), 190–193, 192f  
    in vitro (protocol), 187–189, 188f  
  germlayer explant recombinant culture, 151, 173–174  
  isolating extra-embryonic membranes, 150–151, 167–169, 168f  
  isolating germ cells from genital ridge, 151–152, 152f, 175–177, 176f–177f  
  isolation of embryos, 150, 159–166, 160f–165f  
    early neural-fold-stage dissection, 164f, 165  
    early primitive-streak-stage dissection, 163, 163f  
    early somite-stage dissection, 165–166, 165f  
    implantation-stage dissection, 159–160, 161f  
    late primitive-streak-stage dissection, 163, 164f  
    prestalk-stage dissection, 160, 162, 162f  
  separating postimplantation germ layers, 151, 170–172, 171f  
  visualizing early implantation sites, 150, 150f, 157–158  
Preantral follicle, 30

- Prechordal plate, 53  
 Pregastrula embryo, 45, 47f  
 Pregnancy with manipulated embryos, 94  
 Pregnant mare serum gonadotropin (PMSG), 91, 95, 97, 105–106, 315  
 Preimplantation embryos  
   collection of, 123  
     blastocyst collection (protocol), 146–148, 147f  
     opening abdominal cavity and locating female reproductive organs (protocol), 137–138, 137f, 138f  
     two-cell- to morula-stage embryo collection (protocol), 143–145, 144f–145f  
     zygote collection and removal of cumulus cells (protocol), 139–142, 140f–141f  
 culture of, 109–148  
   considerations for successful, 118–120  
     embryo collection, 118–119  
     medium choice, 118  
     quality control, 119–120  
     reagents, 118–119  
   effect on gene expression, 117–118  
   media, 109–117  
     commercially available, 120t  
     composition of selected media, 111t  
     critical components of, 113–117, 115t  
     for development from zygotes to blastocysts, 112–113  
     history, 109–110  
     for overcoming two-cell block, 109  
     preparation of, 120–121, 126–131, 127t, 128t, 130t  
     microdrop culture method, 110, 119  
     setting up (protocol), 132–133, 133f  
   development of (*see* Development)  
   pipettes for handling, 121–122, 122f, 134–137  
   quality of, 145f  
   transport of, 641, 679–681  
   visualizing early implantation sites, 150, 157–158  
 Primary mouse embryonic fibroblasts (MEFs), 330  
   feeder cell layer preparation from (protocol), 341–342  
   preparation of (protocol), 338–340  
     discussion, 339–340  
     materials, 338–339  
     method, 339  
 Primitive endoderm, 38f, 41, 42f, 490  
 Primordial follicles, 27, 32f  
 Primordial germ cells (PGCs), 49–51, 50f, 51f  
   migration pathway, 50–51, 50f  
 Proestrus, 95t  
 Progenitor cells from the CNS, 334  
   isolation, culture, and differentiation of (protocol), 389–394  
     differentiation, 392  
     discussion, 392  
     dissection, 390–392, 391f  
     materials, 389–390  
     method, 390–392, 390f, 391f  
     recipes, 392–394  
     secondary neurospheres, 392  
     slide coating, 392  
   isolation, culture, and differentiation of radial glial-like neural progenitor cells in adherent cultures (protocol), 395–401  
     coating culture surfaces with AK-cyclo(RGDfC), 396  
     differentiation, 399, 400f  
     dissection, tissue dissociation, and cell seeding, 396–397  
     fixation and immunohistochemistry, 399  
     microinjection of lentivirus into embryos, 308–311  
     microinjection of RNA into zygotes, 306–307  
     microinjection of zygotes, 301–305, 303f  
     microinjection setup, 298–300, 299f, 300f  
     troubleshooting guide, 315–319  
     in vitro-translated RNA for microinjection, preparation of, 292–293  
     YAC DNA purification using PFGE and ultrafiltration, 286–288  
     YAC DNA purification using two-gel electrophoresis procedure, 279–285  
     yeast agarose plug preparation for YAC DNA isolation, 274–278  
     timing of microinjection, 262  
     troubleshooting guide, 315–319  
     zygote preparation, 255–256, 255f  
 Pronuclear transfer  
   overview, 572  
   protocol, 577–582  
     embryo isolation, 578  
     enucleation of zygote, 579–580, 580f  
     materials, 577–578  
     pipette preparation, 578–579, 579f  
     preparation of inactivated Sendai virus, 580  
     pronuclei introduction into enucleated zygotes, 581–582, 581f  
 Proteinase K, 558–560, 564  
 Pseudopregnancy, 201, 202  
   females to serve as embryo recipients, 92–93  
   sterile males for induction of, 93  
 Pullers, 775  
 Pulsed-field gel electrophoresis (PFGE)  
   YAC DNA purification using PFGE and ultrafiltration (protocol), 286–288  
   YAC DNA purification using two-gel electrophoresis procedure (protocol), 279–285  
*puro* (puromycin-resistance gene), 435, 465t  
 Puromycin, 435, 465t, 469t  
 PZD (partial zona dissection) of oocytes, 597
- Q**  
 QIAEX kit/QIAquick kit, 250t  
 Qiagen Endo Free Plasmid Kit, 250t, 464  
 QIAquick kit, 250t  
 Quality control, for embryo culture media, 119–120  
 Quarantine, 86–87, 90
- R**  
 RACE (rapid amplification of cDNA ends), 555  
 Radial glial-like neural progenitor cells (RGLs)  
   isolation, culture, and differentiation of cells in adherent cultures (protocol), 395–401  
     coating culture surfaces with AK-cyclo(RGDfC), 396  
     differentiation, 399, 400f  
     dissection, tissue dissociation, and cell seeding, 396–397  
     fixation and immunohistochemistry, 399
- coating culture surfaces with  
   AK-cyclo(RGDfC), 396  
 differentiation, 399, 400f  
 dissection, tissue dissociation, and cell seeding, 396–397  
 fixation and immunohistochemistry, 399  
 harvesting by trypsinization, 398–399  
 maintenance and propagation of cells, 397–398, 398f, 399f  
 materials, 395–396  
 methods, 396–399, 398f–400f  
 recipes, 400–401  
 Progesterone (Depo-Provera), 204, 233  
 Progesterone receptor, 239, 440  
 Prokaryotic vector sequences, use in transgenes, 242  
 Promoters  
   endogenous, 432  
   exogenous, 432  
   of housekeeping genes, 245  
   tetracycline-regulated (SG-TRE), 440  
   in vector design, 432  
 Promoter trap, 445–446  
 Pronase solution (recipe), 783  
 Pronuclear microinjection  
   detection and analysis of transgene produced from, 552–553  
   embryo donors for, 89–90, 91–92, 92t  
   production of donors, 91–92, 92t  
   strains for, 89–90  
   transgenic mouse production by, 237–319  
   applications of technology, 238–240  
   designing transgenes, 242–247  
   DNA isolation and purification  
     commercial kits for, 249, 250t  
     overview, 247–252  
     preparation of DNA for microinjection, 252–254  
     protocols, 269–291  
   equipment  
     general setup, 258, 259f  
     holding pipettes, 256–257, 256f, 257f  
     injection chambers, 260–262, 260f–262f  
     injection pipettes, 257–258  
     micromanipulators, 259  
     microscope, 258  
     pressure control in holding pipette, 259  
     pressure control in injection pipette, 259–260  
   homologous recombination, 241  
   microinjection technique, 262–263  
   mouse lines, establishment and maintenance of, 263–264  
   mouse strain choice, 254–255  
   overview, 240–242  
   post-injection handling of embryos, 263  
 protocols  
   BAC DNA isolation using NucleoBond system, 271–272  
   BAC DNA purification, 273  
   DNA fragment isolation and purification, 269–270  
   infecting embryos by coculturing with lentivirus, 312–314  
   injection buffer preparation for BAC/YAC DNA, 290  
   injection buffer preparation for standard-sized DNA, 289  
   making holding pipettes, 294–295  
   making injection pipettes, 296–297

- harvesting by trypsinization, 398–399  
 maintenance and propagation of cells, 397–398, 398f, 399f  
 materials, 395–396  
 methods, 396–399, 398f–400f  
 recipes, 400–401
- Radiation**  
 safety procedures, 792  
 waste disposal, 791, 792
- Ramp test**, 514
- Rapid amplification of cDNA ends (RACE)**, 555
- Rapid DNA prep lysis buffer (recipe)**, 482
- Rathke's pouch**, 67, 67f
- Rat serum**, preparation of, 178–179, 180f
- Reactive oxygen species (ROS)**, 116
- Reagents**, quality of, 118–119
- Recipes**  
 acidic Tyrode's solution for removing zona pellucidae from preimplantation embryos, 780  
 activin A stock solution (10 µg/mL), 386  
 alkaline phosphatase (AP) buffer, 717, 752  
 alkaline phosphatase buffer (NTMT), 739  
 alkaline phosphatase buffers, 780  
 alkaline phosphatase staining solution, 780  
 β-mercaptoethanol, 360  
 blocking solution, 718  
 BSA fraction V in PBS (0.1%, w/v), 386  
 BSA fraction V in PBS (0.1%, w/v), 377  
 CaCl<sub>2</sub> (0.1 M), 392  
 chemically defined medium (CDM)+FGF2 and ActA medium for culture in feeder-free conditions, 387  
 chemically defined medium (CDM) for culture in feeder-free conditions, 387  
 Chicago Sky Blue 6B (Pontamine Sky Blue), 781  
 complement, guinea pig serum, 360  
 complete ES cell medium (ES-DMEM), 327t, 430  
 complete ES cell medium for C57BL6/N ES cells, 529  
 complete ES cell medium for 129S6/SvEvTac X C57BL/6N F<sub>1</sub> hybrid ES cells, 530  
 cryoprotectant solution, 1.5 M  
   1,2-propanediol, 650  
 cryoprotective agent with 100 mM  
   L-glutamine (gCPA), 674–675  
 cryoprotective medium (CPM), 669  
 culture medium, 174, 186  
 CZB-HEPES media, 629t–630t  
 CZB media, 629t–630t  
 DEPC-treated water, 697, 705  
 detergent rinse, 749  
 digestion buffer, 559  
 dilution solution, 1 M sucrose, 651  
 dissection medium, 186  
 dissociation medium, 393  
 DMEM/F12 + N2 medium, 360  
 DMSO (1.5 M) cryoprotectant solution, 677  
 EFS42.5c vitrification solution, 663–664  
 EFS20 vitrification solution, 663  
 embryo dilution solution, 657  
 enzyme solution for adult tissue dissociation, 400  
 enzyme solution for neurosphere assay, 393  
 EpiSC + FGF2 medium, 388  
 EpiSC medium, 387  
 feeder-conditioned medium (Feeder-CM), 377  
 FGF2 stock solution (10 µg/mL), 388  
 FGF4 stock solution (1000×, 25 µg/mL), 377  
 fixative for frozen sections, 749  
 fixative for staining intact embryos, 749  
 freezing medium (2×), 477  
 freezing medium for feeder cell culture (2×), 388  
 freezing medium for feeder-free cell culture (1×), 388  
 gelatin (0.1%), 360  
 Gittschier buffer, 486  
 heavy inhibitory solution, 393  
 heparin stock solution (1000×, 1 mg/mL), 377  
 HEPES-buffered DMEM with FBS, 166, 169, 172  
 human tubal fluid medium (HTF), 612, 621  
 human tubal fluid medium with calcium, 613  
 hyaluronidase, 781  
 hybridization mix, 739  
 inhibitors (2i), 10 mM stock, 360  
 KSOM+2i inhibitors (100× stock), 360  
 light inhibitory solution, 394  
 lysis buffer, 484, 565, 570  
 MABT, 739  
 mannitol (0.3 M), 781  
 methylene blue solution, 781  
 MgCl<sub>2</sub> (1 M), 394  
 modified PBS supplemented with glucose, pyruvic acid, BSA, and antibiotics, 613  
 NaCl/ethanol mixture, 482  
 N2B27 medium, 361  
 N2B27 medium with LIF and 2i inhibitors, 361  
 NDS buffer (100%), 276–277  
 NEP basal medium, 394  
 NEP complete medium, 394  
 neurobasal + B27 medium, 360  
 neurobasal medium + PSG, 394  
 neutralization buffer, 734  
 N2 supplement solution (100×), 361  
 N2 supplement (100×) stock solutions, 361  
 NTMT, 739, 780  
 pancreatin/trypsin solution for separating germ and tissue layers, 781  
 PBSMT for immunohistochemistry of whole-mount embryos, 782  
 PBT, 782  
 PCR buffer (1×), 486  
 PCR buffer (10×), 568  
 PCR lysis buffer, 486  
 PCR reaction mixture, 487  
 PCR tissue homogenization buffer (10×), 563  
 phenol:chloroform solution, 782  
 phosphate-buffered saline (PBS), 782–783  
 phosphate-buffered saline (PBS at pH 7.4 with Ca<sup>2+</sup>/Mg<sup>2+</sup>), 740  
 phosphate-buffered saline (PBS at pH 7.4, 10×), 740  
 polyamine mix (1000×), 284  
 polyvinylpyrrolidone (10%–12%), 590  
 pronase solution, 783  
 rapid DNA prep lysis buffer, 482  
 restriction digestion mixture, 482  
 RGL basal medium, 400  
 RGL complete medium, 400  
 saline/EDTA buffer plus glucose, 783  
 salts (10×), 734  
 scaleA2, 754  
 70cond + F4H medium, 378  
 70cond medium, 377  
 solution I for YAC DNA agarose plug formation, 277  
 solution II for YAC DNA agarose plug formation, 277  
 solution III for YAC DNA agarose plug formation, 277  
 solution IV for YAC DNA agarose plug formation, 277  
 S-PB1 dilution solutions, 664  
 SSC (20×), 783  
 SSPE (20×), 783  
 staining solution, 750  
 TAE (Tris-acetate/EDTA) buffer (1×), 783  
 TBE buffer (10×), 285  
 TBST, 718  
 TE (Tris-EDTA) buffer, 783  
 TES buffer, 697  
 transcription buffer (5×), 734  
 trypsin-EDTA solution, 784  
 trypsin (0.25%) in Tris-saline, 784  
 trypsin versene phosphate (TVP) buffer, 362  
 TS + F4H medium (1×), 378  
 TS medium, 378  
 TYH medium supplemented with MBCD, 614  
 Tyrode Ringer's saline (pH 7.6–7.7), Ca<sup>2+</sup>/Mg<sup>2+</sup> free, 784  
 YAC equilibrium buffer, 285
- Recombinases**, 437–439. *See also* Site-specific recombinases; *specific recombinases*
- Recombination-mediated cassette exchange (RMCE)**, 448–449
- Rederivation of mouse strains**, 637–640
- Refrigerators and freezers**, 769
- Reichert's membrane**, 45, 167, 168f, 169
- Replica-plating**, 463, 463f
- Reporters**, 436–437, 437f. *See also specific genes*  
 conditional reporter lines, 15  
 to detect recombinase action, 440  
 fluorescent proteins, 436–437, 437f  
*hPLAP* (human placental alkaline phosphatase) gene, 437  
*lacZ* gene, 16, 17, 436  
 in transgene design, 244, 436–437, 437f
- Reproductive biology statistics**, 4t
- Reprogramming of somatic cells to induced pluripotent stem (iPS) cells**  
 overview, 406–407  
 protocols  
   integrating *piggyBac* transposon transgenes into mouse fibroblasts by electroporation, 424–425  
   integrating *piggyBac* transposon transgenes into mouse fibroblasts using chemical methods, 426–427  
   reprogramming mouse fibroblasts with *piggyBac* transposons, 428–430, 429f  
   vectors for, 407–408, 407f
- Resources**, WWW, 785–787
- Restriction digestion mixture (recipe)**, 482
- Restriction enzymes**, 462, 462t
- Restriction fragment length polymorphism (RFLP) studies**, 4
- Retroviruses**, 462
- RGL basal medium (recipe)**, 400

- RGL complete medium (recipe), 400  
RGLs. *See* Radial glial-like neural progenitor cells  
RI strains, 8–9, 8t  
RMCE (recombination-mediated cassette exchange), 448–449
- RNA  
concentration measurement, 293  
isolating total RNA from embryos or fetal tissues, 684, 695–697, 696t  
microinjection of RNA into zygotes (protocol), 306–307  
in vitro-translated RNA for microinjection, preparation of (protocol), 292–293
- RNA interference, 15
- RNA probes  
in situ hybridization of embryo and tissue sections with radiolabeled RNA probes (protocol), 728–732  
autoradiography, 733–734  
hybridization, 731  
materials, 728–729  
posthybridization washing, 731–732  
prehybridization, 730–731  
probe preparation, 729–730  
in situ hybridization of whole-mount embryos with nonradiolabeled RNA probes (protocol), 735–740  
discussion, 738–739  
embryo preparation, 736  
histochemistry, 738  
materials, 735–736  
post-antibody washes, 738  
posthybridization washes, 738  
pretreatment and hybridization, 737  
probe preparation, 736–737  
recipes, 739–740
- RNase, avoiding contamination with, 687
- Robertson, Elizabeth J., 15
- Robertsonian chromosomes, 9
- Roller culture of postimplantation embryos  
overview, 153  
protocol, 178–181, 179f–180f  
culturing embryo, 180–181  
embryo preparation, 180–181  
gas mixture for embryos, 179f  
materials, 178  
serum preparation, 178–179, 180f
- ROS (reactive oxygen species), 116
- Rosa26 locus, 245, 458
- Roux, Wilhelm, 12
- rox sequence, 438
- RU-486, 440
- Ruddle, Frank, 14
- Russell, Liane, 10
- S**
- Safety, 789–794  
biological safety procedures, 792–793  
general cautions, 789–791  
institutional safety office, 789  
material safety data sheets (MSDSs), 789  
properties of common hazardous chemicals, 793–794  
radioactive safety procedures, 792  
waste disposal, 791
- Saline/EDTA buffer plus glucose (recipe), 783
- Salts  
10× (recipe), 734  
as DNA prep contaminant, 248
- Sato, Gordon, 13
- Scale, 753
- ScaleA2 (recipe), 754
- Sclerotome, 57, 57f
- SCNT. *See* Somatic cell nuclear transfer (SCNT)
- Scrotal method for vasectomy, 209f, 210
- Sectioning. *See* Tissue sections
- Selectable markers, 434–436  
dt (diphtheria toxin), 436  
hprt (hypoxanthine phosphoribosyltransferase), 435, 436  
HSV-tk (herpes simplex virus-1 thymidine kinase gene), 436  
hygro (hygromycin-resistance gene), 435  
negative, 436  
neo (neomycin phosphotransferase), 434–435  
positive, 434–435  
puro (puromycin-resistance gene), 435
- Selection methods, 465, 465t
- Seminiferous tubules, 224, 404, 405  
microinjection of spermatogonial stem cells into, 420–422, 421f
- Sendai virus, cell fusion induced by, 572, 580–581, 581f
- Sensory system, development stages, 26t–27t
- Sequential media (G1 and G2), 111t, 113
- Serum batches, testing (protocol), 348–349
- Serum preparation from rats, 178–179, 180f
- 70cond + F4H medium (recipe), 378
- 70cond medium (recipe), 377
- Sex chromosomes in chimeras, 493, 500
- Sex determination and differentiation, 60–61
- Sharpening tungsten needles by electrolysis, 160f
- short ear locus, 10
- Silica matrix columns, for contaminant removal in DNA prep, 249
- Site-specific recombinases, 437–439. *See also specific recombinases*  
Cre recombinase, 437–438  
detection and analysis of ES cell-mediated gene/genome alterations induced by, 554  
Dre recombinase, 438  
φC31 integrase, 439  
Flp recombinase, 439  
in recombination-mediated cassette exchange (RMCE), 448–449  
reporters to detect, 440
- Site-specific recombination-mediated insertions, 448–449
- Skeleton, staining of, 690f, 693–694
- Skin preparation surgical procedures, 196
- Škreb, Nikola, 13
- Sleeping Beauty*, 17, 241, 407
- Smithies, Oliver, 15, 18
- Snell, George, 7–8
- Solter, Davor, 13
- Solution I for YAC DNA agarose plug formation (recipe), 277
- Solution II for YAC DNA agarose plug formation (recipe), 277
- Solution III for YAC DNA agarose plug formation (recipe), 277
- Solution IV for YAC DNA agarose plug formation (recipe), 277
- Solvents, safe handling of, 790, 793
- Somatic cell nuclear transfer (SCNT), 403, 406  
cloning mice (protocol), 583–590  
Honolulu method, 572
- Somites, 55–58, 56f, 57f
- sonic hedgehog gene, 61, 65
- Southern blot analysis  
for copy number determination, 553  
for detection and analysis of ES cell-mediated gene/genome alterations, 554  
DNA preparation from cultured cells for (protocol), 567–568  
for genotyping, 552–553  
identification of homozygous transgenic mice/embryos using flanking probe, 556
- SOX2 transcription factor and gene, 406, 407
- S-PB1 dilution solutions (recipes), 664
- Specific pathogen-free (SPF) facilities, 86, 87, 89  
rederivation of mouse strains, 637–639
- Spectral karyotyping (SKY), 332
- Spemann, Hans, 10, 12
- Sperm  
collection for in vitro fertilization, 606, 609  
cryopreservation  
overview, 635–636  
using cryoprotectant containing L-glutamine (protocol), 671–675, 672f–674f  
using cryoprotectant containing monothioglycerol (MTG) (protocol), 665–670  
cryopreservation of, 597  
intracytoplasmic sperm injection (ICSI), 598, 623–630, 626f, 627f  
preparation of cryorecovered for IVE, 607, 609  
transport of samples, 640
- Spermatogenesis, 31f, 32–34, 404
- Spermatogonial stem cells (SSCs)  
overview, 404–406  
culture of, 404–405  
transplantation of, 405–406, 405f  
protocols  
cryopreserving and thawing spermatogonial stem cells, 416–417  
culture and expansion of primary undifferentiated spermatogonial stem cells, 413–415  
establishment of primary culture, 413–414, 415f  
expansion of primary undifferentiated culture, 414–415  
materials, 413  
isolation of the spermatogonial stem cell-containing fraction from testes, 410–412  
materials, 410–411  
percoll isolation, 412  
testes disaggregation, 411  
Thy1<sup>+</sup> selection by magnetic-activated cell sorting, 412  
spermatogonial stem cell transplantation to the testis, 418–423  
busulfan treatment of adult recipients, 419  
discussion, 422  
injection of cells into recipient tissue, 420–422, 421f  
materials, 418–419, 419f  
preparation of cells for transplantation, 420  
preparation of injection pipette and injector, 419–420



- Sperm count depression after mating, 91  
 Splenectomy, 228  
 Splice acceptor, 432  
 Splice donor, 432  
 Splinkerette PCR, 555  
 Sry gene, 60  
 SSC (saline sodium citrate) (20×) (recipe), 783  
 SSCs. *See* Spermatogonial stem cells  
 SSEA1, for testing pluripotency and germline competence of induced pluripotent stem (iPS) cells, 408, 408f  
 SSPE (20×) (recipe), 783  
 Staining for alkaline phosphatase activity, 689f, 691, 751–752  
 Staining for β-galactosidase activity, 689f, 691  
 Staining frozen sections (protocol), 748–750  
 Staining solution (recipe), 750  
 Staining whole embryos and sections for β-galactosidase (LacZ) activity (protocol), 745–747  
 Static culture of postimplantation embryos, 153–154, 182–186  
   culture medium preparation, 183  
   for imaging, 154, 184–186, 185f  
   overview, 153–154  
   setup, 183  
 STAT3 pathway, 324  
 Steel pigmentation mutant, 9  
 Stem cells. *See also* Embryonic stem (ES) cell(s); Induced pluripotent stem cells; Spermatogonial stem cells; Trophoblast stem cells  
   germline-competent derived from adult mice, 403–430  
   subcutaneous injection of pluripotent stem cells (protocol), 225–226  
   transplantation under kidney capsule (protocol), 222–224, 223f  
 Stereomicroscopes, 771  
 Stevens, Leroy, 13  
 STO fibroblasts, 330–331  
   feeder cell layer preparation from (protocol), 341–342  
 Strain distribution pattern, 8  
 Strains  
   for chimera formation, 491–493, 494f–499f, 494t–500t  
   creation of inbred, 4, 7–9  
   for embryo donors  
     for generation of ES cell chimeras, 90  
     for pronuclear microinjection, 89–90  
   genealogy of commonly used inbred strains, 5f–6f  
   origin of the laboratory mouse, 4, 5f–7f  
   for pronuclear microinjection, 254–255  
   rederivation of, 637–640  
   for superovulation, 96, 96t  
   vendors of, 768  
 Strong, Leonell C., 7  
 Strontium-induced parthenogenesis, 572, 576, 576f  
 Subcutaneous injection of pluripotent stem cells (protocol), 225–226  
 Subzonal insemination, 598  
 Superovulation induction, 95–97, 96t  
   age and weight, 96–97, 96t  
   gonadotropin  
     dose, 96t, 97  
     preparation of, 105  
     time of administration, 97  
   performance of stud males, 97  
   protocol, 105–106  
   strain background, 96, 96t  
 Surgical instruments, 776–777  
   care and cleaning of, 196  
   commercial sources of, 196  
   sterilization of, 196  
 Surgical procedures, 195–235  
   analgesia, 198–199, 199t  
   anesthesia, 197–198, 198t  
   caesarean section, 203, 220–221  
   castration, 204, 234–235  
   embryo transfer, 199–203  
     background, 199–200  
     nonsurgical embryo transfer (NSET), 200  
   number of embryos to transfer, 200–201  
   protocols  
     oviduct transfer of embryos, 211–215, 212f, 213f  
     uterine transfer of embryos, 216–219, 218f  
     recipient females, 201–202  
     technical aspects, 202–203, 202f  
     unilateral and bilateral transfers, 201  
     zona pellucida role in, 200  
   general guidelines for, 196–197  
   closure, 197  
   incisions, 196  
   instrument care and sterilization, 196  
   post-surgical care, 197  
   skin preparation, 196  
   isolating inner cell mass of blastocysts (protocol), 549–550, 549f  
   location of embryo manipulation laboratory, 87–88  
   ovariectomy, 203–204, 232–233  
   ovary transplantation, 596, 601–603, 602f  
   protocols  
     blood collection by tail bleeding, 230–231  
     caesarean section and fostering, 220–221  
     castration, 234–235  
     opening abdominal cavity and locating female reproductive organs, 137–138, 137f, 138f  
     ovariectomy, 232–233  
     oviduct transfer of embryos, 211–215, 212f, 213f  
     subcutaneous injection of pluripotent stem cells, 225–226  
     tissue biopsy, 227–229  
     tissue transplantation under kidney capsule, 222–224, 223f  
     tribromoethanol preparation, 206–207  
     uterine transfer of embryos, 216–219, 218f  
     vasectomy, 208–210, 209f  
   tissue biopsy, 203, 227–229  
   tissue transplantation  
     overview, 203  
     protocols  
       subcutaneous injection of pluripotent stem cells, 225–226  
       transplantation under kidney capsule, 222–224, 223f  
       vasectomy, 199, 208–210, 209f  
 Suspension culture, differentiating ES cells into embryoid bodies (protocol), 368–369  
 Systematic mutagenesis and phenotyping, 15–18
- ## T
- Tables, vibration-free, 773  
 TAE (Tris-acetate/EDTA) buffer (1×) (recipe), 783  
 Tail bleeding, blood collection by (protocol), 230–231  
 Tail bud, 54  
 Tail tip excision, 100  
 Takahashi, Kazutoshi, 14, 406  
 TALE nucleases (TALEN), 1, 434, 434f  
 TALE (transcription activator-like effector) peptides, 434  
 Tamoxifen, 204  
 Tamoxifen-inducible system, 239, 440  
 Tarkowski, Andrzej, 490  
 Tarkowski, Kristof, 12  
 Tattooing, 98  
 Taylor, Benjamin A., 8  
 TBE (Tris/borate/EDTA) buffer (10×) (recipe), 285  
 TBST (Tris-buffered saline + Tween 20) (recipe), 718  
 T4 DNA polymerase, 739  
 TE (Tris-EDTA) buffer (recipe), 783  
 Temperature, for embryo culture, 117  
 Temperature controllers for heating and cooling microscope stages, 772  
 Teratoma  
   from epiblast stem cell (EpiSC), 44  
   subcutaneous injection of pluripotent stem cells (protocol), 225–226  
 TES buffer (recipe), 697  
 TESP (3-triethoxysilylpropylamine), 685  
 Test breeding, for identification of homozygous transgenic mice/embryos, 556  
 Testis  
   capsule, tissue transplantation under, 224  
   isolation of the spermatogonial stem cell-containing fraction from (protocol), 410–412  
   materials, 410–411  
   percoll isolation, 412  
   testes disaggregation, 411  
   Thy1<sup>+</sup> selection by magnetic-activated cell sorting, 412  
   spermatogonial stem cell transplantation to (protocol), 418–423  
   busulfan treatment of adult recipients, 419  
   discussion, 422  
   injection of cells into recipient tissue, 420–422, 421f  
   materials, 418–419, 419f  
   preparation of cells for transplantation, 420  
   preparation of injection pipette and injector, 419–420  
 tetO, 239, 407, 407f, 440  
 Tetracycline-inducible system, 239, 440  
   described, 440  
   for reprogramming of somatic cells to induced pluripotent stem (iPS) cells, 407, 407f, 428–430  
 Tetracycline repressor (TetR), 239, 440  
 Tetracycline transactivators, 458  
 Tetraploid embryo  
   chimeras types  
     diploid embryo ↔ tetraploid embryo, 501t, 502, 544–545  
     ES cells ↔ tetraploid embryo, 501t, 502–503, 545–546, 546f

- Tetraploid embryo (*Continued*)  
  production of (protocol), 539–541, 540f, 540t  
*T* gene, 10–11  
TGF- $\alpha$  (transforming growth factor- $\alpha$ ), 117  
TGF- $\beta$  (transforming growth factor- $\beta$ ), 60  
Thawing  
  embryos, 649  
  ES cells, 346–347  
    from a 96-well plate, 478–479  
  sperm, 668, 673–674  
  spermatogonial stem cells, 416–417  
Thy1<sup>+</sup> selection by magnetic-activated cell sorting, 412  
Tissue adhesive, 197  
Tissue biopsy  
  ear punching, 98  
  overview, 203  
  protocol, 227–229  
    materials, 227  
    nephrectomy, 228–229  
    partial hepatectomy, 227–228  
    splenectomy, 228  
  tail tip excision, 100  
Tissue fixation  
  fixation and paraffin embedding for GFP visualization (protocol), 757–758  
  handling blastocysts for fixation (protocol), 700  
  overview, 685  
  solution preparation (Bouin's fixative) (protocol), 699  
  solution preparation (paraformaldehyde 4%) (protocol), 698  
Tissue-nonspecific alkaline phosphatase (TNAP), 49  
Tissue sections  
  sectioning embryos after whole-mount in situ hybridization (protocol), 742–743  
  staining frozen sections (protocol), 748–750  
  transferring sections onto water droplets (protocol), 708  
Tissue-specific regulatory sequences, 244–245  
Tissue transplantation  
  overview, 203  
  protocols  
    subcutaneous injection of pluripotent stem cells, 225–226  
    transplantation under kidney capsule, 222–224, 223f  
    under testis capsule, 224  
TNAP (tissue-nonspecific alkaline phosphatase), 49  
Toe clipping, 98  
*Tol2*, 240  
Toxic compounds, 794  
Transcription activator-like effector (TALE) peptides, 434  
Transcription buffer (5 $\times$ ) (recipe), 734  
Transcription factors, Yamanaka, 406, 407  
Transfection methods, 462–463  
Transforming growth factor- $\alpha$  (TGF- $\alpha$ ), 117  
Transforming growth factor- $\beta$  (TGF- $\beta$ ), 60  
Transgenes  
  copy number determination, 553  
  designing, 242–247  
    coinjection of multiple transgenes, 243  
    distinguishing expression of transgenes and endogenous genes, 243–244  
    efficiency of gene transfer, factors affecting, 245–247  
    buffer used, 246  
    concentration of DNA, 246  
    linear or circular form of the DNA, 246  
    purity of DNA, 246  
    size of DNA construct, 245, 246  
  expression of cDNAs, 245  
  housekeeping gene use, 245  
  introns, role of, 245  
  large constructs, issues with, 245–247  
  length of construct, 242–243  
  prokaryotic vector sequences, effects of, 242  
  reporter genes, 244  
  tissue-specific regulatory sequences, 244–245  
  ubiquitous expression, 245  
  detection and analysis of transgene produced by pronuclear injection, 552–553  
Transgenic mice  
  applications of technology, 238–240  
  designing transgenes, 242–247  
  coinjection of multiple transgenes, 243  
  distinguishing expression of transgenes and endogenous genes, 243–244  
  efficiency of gene transfer, factors affecting, 245–247  
  expression of cDNAs, 245  
  housekeeping gene use, 245  
  introns, role of, 245  
  large constructs, issues with, 245–247  
  length of construct, 242–243  
  prokaryotic vector sequences, effects of, 242  
  reporter genes, 244  
  tissue-specific regulatory sequences, 244–245  
  ubiquitous expression, 245  
  genotyping, 551–570  
  identification of homozygous, 555–557  
  production by pronuclear microinjection, 237–319  
    applications of technology, 238–240  
    designing transgenes, 242–247  
    DNA isolation and purification  
      commercial kits for, 249, 250t  
      overview, 247–252  
      preparation of DNA for microinjection, 252–254  
      protocols, 269–291  
  equipment  
    general setup, 258, 259f  
    holding pipettes, 256–257, 256f, 257f  
    injection chambers, 260–262, 260f–262f  
    injection pipettes, 257–258  
    micromanipulators, 259  
    microscope, 258  
    pressure control in holding pipette, 259  
    pressure control in injecting pipette, 259–260  
  homologous recombination, 241  
  microinjection technique, 262–263  
  mouse lines, establishment and maintenance of, 263–264  
  mouse strain choice, 254–255  
  overview, 240–242  
  post-injection handling of embryos, 263  
  protocols  
    BAC DNA isolation using NucleoBond system, 271–272  
    BAC DNA purification, 273  
    DNA fragment isolation and purification, 269–270  
    infecting embryos by coculturing with lentivirus, 312–314  
    injection buffer preparation for BAC/YAC DNA, 290  
    injection buffer preparation for standard-sized DNA, 289  
    making holding pipettes, 294–295  
    making injection pipettes, 296–297  
    microinjection of lentivirus into embryos, 308–311  
    microinjection of RNA into zygotes, 306–307  
    microinjection of zygotes, 301–305, 303f  
    microinjection setup, 298–300, 299f, 300f  
    troubleshooting guide, 315–319  
  in vitro-translated RNA for microinjection, preparation of, 292–293  
  YAC DNA purification using PFGE and ultrafiltration, 286–288  
  YAC DNA purification using two-gel electrophoresis procedure, 279–285  
  yeast agarose plug preparation for YAC DNA isolation, 274–278  
Translocations  
  in ES cell lines, 332–333  
  induced site-specific, 450–451  
Transport of samples  
  of cryopreserved embryos and sperm, 640  
  of dissected oviducts, 641  
  of live freshly collected samples, 640–641  
  of preimplantation embryos, 641, 679–681  
  packaging, 679–680  
  receiving embryos, 680  
Transposon-based reprogramming, vectors for, 407–408, 407f  
Tribromoethanol  
  anesthesia with, 197–198  
  preparation protocol, 206–207  
  materials, 206  
  using concentrated stock, 206–207  
  working solution (1.25%), 207  
Trichostatin A (TSA), 573  
Tris/borate/EDTA buffer. *See* TBE buffer  
Tris-buffered saline + Tween 20. *See* TBST buffer  
Trisomies in ES cells, 332–333  
Trophoblast, 73  
Trophoblast stem (TS) cells, 333–334  
  derivation and culture of TS cell lines (protocol), 373–378  
  discussion, 376  
  materials, 373–374  
  method, 374–376, 375f  
  recipes, 376–378  
Trophectoderm, 36–37, 37f, 38f, 39, 41, 42f, 490, 549–550  
Trypsin, 330, 528, 533  
  in chromosomal banding, 332  
  harvesting RGLs by trypsinization, 398–399  
Trypsin-EDTA solution (recipe), 784  
Trypsin (0.25%) in Tris-saline (recipe), 784  
Trypsin versene phosphate (TVP) buffer (recipe), 362  
TSA (trichostatin A), 573  
TS cells. *See* Trophoblast stem (TS) cells

- TS + F4H medium (1×) (recipe), 378  
 TS medium (recipe), 378  
 Tungsten needles, sharpening by electrolysis, 160f  
 Turning, embryonic, 54, 55f  
 2A peptide, 243–244, 433  
 Two-cell block, culture media for overcoming, 109  
 2i (PD0325901 and CHIR99021), derivation of ES cell lines using (protocol), 355–362  
 embryo appearance, 358f  
 expansion of ES cell lines, 359, 359f  
 inner cell mass isolation by immunosurgery, 357–359  
 materials, 355–356  
 preimplantation epiblast preparation, 356–357  
 recipes, 360–362  
 summary diagram, 357f  
 TYH medium supplemented with methyl- $\beta$ -cyclodextrin (MBCD) (recipe), 614  
 Tyrode Ringer's saline (pH 7.6–7.7),  $\text{Ca}^{2+}/\text{Mg}^{2+}$  free (recipe), 784
- U**  
 Ultrafiltration, in DNA purification, 287–288, 287f  
 Ultrasonicators, 790–791  
 Urogenital system, development stages, 26t–27t  
 Uterine transfer of embryos (protocol), 216–219, 218f  
 materials, 216  
 method, 217–219, 218f  
 Uterus  
 blastocyst collection from, 146–148, 147f  
 blastocyst implantation, 37, 39, 40f, 149  
 electroporation of postimplantation embryos in utero (protocol), 190–193, 192f  
 isolation of postimplantation embryos from, 150, 159–166, 160f–165f  
 visualizing early implantation sites, 150, 150f, 157–158
- V**  
 Vaginal plug, 201  
 appearance of, 104f  
 checking for, 103  
 Vaginal smear, estrous cycle stage determination by (protocol), 617–618, 618f  
 Vasculature visualization by India ink injection, 690f, 694, 763–764  
 Vasectomy  
 overview, 199  
 protocol, 208–210, 209f  
 abdominal method, 208–210, 209f  
 materials, 208  
 scrotal method, 209f, 210  
 pseudopregnancy induced by vasectomized mice, 93  
 Vector design, 17, 431–460  
 ES cell mediated transgenesis, 440–442  
 examples, 441–442  
 overview, 440–441  
 in vitro screen for widespread transgene expression (protocol), 455–457  
 ES cells homozygous for specific mutations, 451–452  
 gene and promoter trap, 445–446  
 gene targeting, 442–445  
 glossary of elements, 432–440  
 genomic and gene elements, 432–434  
 arms of homology, 433–434, 434f  
 exon, 432  
 internal ribosomal entry site (IRES), 432  
 intron, 432  
 polyadenylation signal, 432  
 promoters, 432  
 splice donor and splice acceptor, 432  
 2A peptide sequence, 433  
 inducible systems, 439–440  
 tamoxifen, 440  
 tetracycline, 440  
 reporters, 436–437, 437f  
 to detect recombinase action, 440  
 fluorescent proteins, 436–437, 437f  
 hPLAP (human placental alkaline phosphatase) gene, 437  
 lacZ gene, 436  
 selectable markers, 434–436  
 dt (diphtheria toxin), 436  
 hprt (hypoxanthine phosphoribosyltransferase), 435, 436  
 HSV-tk (herpes simplex virus-1 thymidine kinase gene), 436  
 hygro (hygromycin-resistance gene), 435  
 negative, 436  
 neo (neomycin phosphotransferase), 434–435  
 positive, 434–435  
 puro (puromycin-resistance gene), 435  
 site-specific recombinases, 437–439  
 Cre recombinase, 437–438  
 Dre recombinase, 438  
 $\Phi\text{C31}$  integrase, 439  
 Flp recombinase, 439  
 reporters to detect, 440  
 tetracycline-inducible system, 440  
 homologous recombination, replacement by, 446–448  
 hit and run example, 447  
 piggyBac transposon example, 448  
 tag and exchange example, 447  
 induced site-specific chromosomal aberrations, 449–451  
 deletions, 450  
 inversions, 450  
 translocations, 450–451  
 protocols, 455–460  
 in vitro screen for widespread transgene expression, 455–457  
 in vitro screen to identify silent but activatable integration sites for tet-inducible transgene, 458–460  
 site-specific recombination-mediated insertions, 448–449  
 vectors for reprogramming of somatic cells to induced pluripotent stem (iPS) cells, 407–408, 407f  
 Ventral ectodermal ridge, 54  
 Vibration-free tables, 773  
 Vibratome, cutting thick sections using (protocol), 709–711  
 fixed specimens, 710–711  
 live specimens, 712–713  
 materials, 709, 712  
 Visceral endoderm, 68–71, 69f–71f, 73  
 Visceral yolk sac  
 development, 68–69, 69f, 72  
 isolation of, 151, 168f, 169  
 Visualization techniques, 683–766  
 for cells using fluorescent proteins, 690f, 692–693  
 compound, confocal, and deconvolution microscope use, 693  
 dissecting microscope use, 692–693  
 macroscopic observation, 692  
 observation in living cells, 693  
 for gene products, 684–691  
 cutting sections, 685  
 dehydrating in alcohol, 685  
 dewaxing and rehydrating sections, 686  
 embedding in wax, 685  
 glass slide and coverslip preparation, 685  
 immunohistochemistry  
 of embryo sections, 686  
 fluorescent analysis, 686  
 general techniques for, 684–686  
 of whole-mount embryos, 686, 689f  
 isolating total RNA from mouse embryos or fetal tissues, 684  
 in situ hybridization  
 avoiding contamination with RNase, 687  
 of embryo and tissue sections with RNA probes, 686–687  
 general techniques for, 684–686  
 of whole-mount embryos with RNA probes, 687–688, 689f, 691  
 staining for alkaline phosphatase activity, 689f, 691  
 staining for  $\beta$ -galactosidase activity, 689f, 691  
 tissue fixation, 685  
 protocols, 695–766  
 Alcian Blue/Alizarin Red staining of cartilage and bone, 760–761  
 Alcian Blue staining of fetal cartilaginous skeleton, 759  
 Alizarin Red staining of postnatal bone, 762  
 autoradiography, 733–734  
 cutting thick sections using a vibratome, 709–711  
 fixed specimens, 710–711  
 live specimens, 712–713  
 materials, 709, 712  
 cutting wax sections using a microtome, 706–707  
 DAPI staining of whole-mount embryos or fetal organs, 722–723, 723f  
 dewaxing and rehydrating sections before in situ hybridization or staining, 714  
 embedding samples in wax, 701–702  
 embedding small embryos in wax, 703  
 fixation and paraffin embedding for GFP visualization, 757–758  
 fluorescent protein observation in living cells, 755  
 glass slide and coverslip preparation for in situ hybridization, 704–705  
 great vessel visualization by plastic coating, 765–766  
 handling blastocysts for fixation, 700

- Visualization techniques (*Continued*)  
  imaging embryos after whole-mount in situ hybridization, 741  
  immunofluorescent staining of whole-mount embryos, 724–725  
  immunohistochemistry of embryo sections, 715–718  
  immunohistochemistry of whole-mount embryos, 719–721  
  Intracellular observation of fluorescent proteins in fixed cells, 756  
  isolating total RNA from embryos or fetal tissues, 695–697, 696t  
  mounting embryos for microscopic observation and imaging, 726, 727f  
  sectioning embryos after whole-mount in situ hybridization, 742–743  
  in situ hybridization of embryo and tissue sections with radiolabeled RNA probes, 728–732  
    autoradiography, 733–734  
    hybridization, 731  
    materials, 728–729  
    posthybridization washing, 731–732  
    prehybridization, 730–731  
    probe preparation, 729–730  
  in situ hybridization of whole-mount embryos with nonradiolabeled RNA probes, 735–740  
    discussion, 738–739  
    embryo preparation, 736  
    histochemistry, 738  
    materials, 735–736  
    post-antibody washes, 738  
    posthybridization washes, 738  
    pretreatment and hybridization, 737  
    probe preparation, 736–737  
    recipes, 739–740  
  staining for alkaline phosphatase activity, 751–752  
  staining frozen sections, 748–750  
  staining whole embryos and sections for  $\beta$ -galactosidase (LacZ) activity, 745–747  
  tissue fixation solution preparation (Bouin's fixative), 699  
  tissue fixation solution preparation (paraformaldehyde 4%), 698  
  transferring sections onto water droplets, 708  
  vasculature visualization by India ink injection, 763–764
- visualization of fluorescent protein expression in whole-mount postimplantation-stage embryos, 753–754  
  wax sections, 744  
  for tissues and organ systems, 693–694  
  fetal great blood vessels by plastic casting, 690f, 694  
  fetal vasculature by India ink injection, 690f, 694  
  skeleton using histochemical stains, 690f, 693–694
- Vitrification, 634, 656  
  embryo cryopreservation by high-osmolality vitrification (protocol), 658–664, 660f, 661f
- Volume of embryo culture, 117
- W**  
Waddington, Conrad, 12  
Wakayama, Teruhiko, 13  
Waltzing mice, 4  
Waste, disposal of, 791, 792  
*waved-1* and *waved-2* hair mutants, 9  
Wax  
  cutting wax sections using a microtome (protocol), 706–707  
  dewaxing and rehydrating sections, 686, 714  
  embedding in  
    fixation and paraffin embedding for GFP visualization (protocol), 757–758  
    overview, 685  
    protocol, 701–702  
    small embryos (protocol), 703  
Weaning, 88  
Whitten, Wesley, 12  
Whitten effect, 95  
Wolffian duct, 60  
Wound clips, 197  
WWW resources, 785–787
- X**  
XEN cells. *See* Extra-embryonic endoderm stem (XEN) cells  
X-gal, 745–746  
X-inactivation, 8, 10
- Xylazine for anesthesia, 198, 198t  
XY sex determination, 60–61
- Y**  
YAC equilibrium buffer (recipe), 285  
Yamanaka, Shinya, 14, 18, 406  
Yamanaka factors, 406, 407  
Yanagimachi, Ryuzo, 13  
Y chromosome, 4  
Yeast agarose plug preparation for YAC DNA isolation (protocol), 274–278  
Yeast artificial chromosomes (YACs), 238, 242–243, 246–247  
  DNA purification using PFGE and ultrafiltration (protocol), 286–288  
  DNA purification using two-gel electrophoresis procedure (protocol), 279–285  
  injection buffer preparation for BAC/YAC DNA (protocol), 290  
  yeast agarose plug preparation for YAC DNA isolation (protocol), 274–278  
Yolk sac  
  development, 68–69, 69f  
  isolation of, 151
- Z**  
Zinc-finger nucleases (ZFNs), 241, 433, 434f  
Zona pellucida, 30, 32f, 34, 37f  
  removal of, 536–538, 537f  
Zona reaction, 34  
Zone of polarizing activity (ZPA), 61  
ZP3 glycoprotein, 34  
Zygotes  
  cleavage, 35  
  collection of (protocol), 139–142, 140f–141f  
  cumulus cell removal with hyaluronidase, 141  
  enucleation of, 579–580, 580f  
  microinjection of zygotes  
    overview, 262–263  
    protocol, 301–305, 303f  
    discussion, 304–305  
    materials, 301  
    method, 301–304, 303f  
    with RNA (protocol), 306–307  
  morphology of, 255f